

Received 2023-04-05

Revised 2023-04-14

Accepted 2023-08-19

The Role of Nursing Care in the Management of Patients with Traumatic Subarachnoid Hemorrhage

Qun Miao ^{1✉}, Yan Yan ¹, Mengjie Zhou ¹, Xueqi Sun ¹¹ Department of Neurosurgery, Funan County People's Hospital, Fuyang, Anhui 236300, China

Abstract

Traumatic subarachnoid hemorrhage (tSAH) is a critical condition that requires comprehensive management to optimize patient outcomes. Nursing care plays a key role in the overall management of patients with tSAH via various aspects of care, including neurological assessment, monitoring, intervention, and education. In this review, we aim to evaluate the significant contributions of nursing care in managing patients with tSAH. Nurses perform initial neurological assessments, including the glasgow coma scale, pupil reactivity, vital signs, and sensory-motor evaluations. These assessments provide valuable information for early identification of deteriorating neurological status and prompt intervention. Additionally, nurses closely monitor intracranial pressure (ICP), cerebral perfusion pressure, and other hemodynamic parameters, assisting in the prevention and timely detection of secondary brain injury. For example, some strategies to manage ICP include elevating the head of the bed, maintaining adequate oxygenation and ventilation, administering proper medications, and ensuring fluid and electrolyte balance. Also, through careful monitoring, early recognition, and appropriate preventive measures, nursing care could prevent complications, including infections, deep vein thrombosis, and pressure ulcers. Furthermore, nursing care extends beyond physical management and encompasses psychosocial support for patients and their families. Nurses establish therapeutic relationships, providing emotional support, education, and counseling to alleviate anxiety, address concerns, and facilitate coping mechanisms. Education regarding medication management, lifestyle modifications, and the importance of regular follow-up enhances patient compliance and promotes long-term recovery. [GMJ.2023;12:e3013] DOI:[10.31661/gmj.v12i.3013](https://doi.org/10.31661/gmj.v12i.3013)

Keywords: Traumatic Subarachnoid Hemorrhage; Nursing Care; Brain Injuries; Intracranial Pressure; Emotional Support

Introduction

Traumatic subarachnoid hemorrhage (tSAH) is a critical condition characterized by bleeding into the subarachnoid space following a traumatic brain injury [1]. It poses a significant challenge for healthcare pro-

fessionals, particularly nurses, which play a crucial role in managing patients with tSAH [2]. The complex nature of tSAH demands a comprehensive nursing care approach that includes assessment, intervention, education, and ongoing support [3, 4]. Hence, this study aimed to short review the valuable role of

GMJ

Copyright© 2023, Galen Medical Journal.
This is an open-access article distributed
under the terms of the Creative Commons
Attribution 4.0 International License
(<http://creativecommons.org/licenses/by/4.0/>)
Email:info@gmj.ir



✉ Correspondence to:

Qun Miao, Department of Neurosurgery, Funan County
People's Hospital, No. 36, Santa Road, Funan County,
Fuyang, Anhui 236300, China
Telephone Number: 0558-6712834
Email Address: jyq4568@163.com

nursing care in managing patients with tSAH, highlighting the essential responsibilities of nurses and their impact on patient outcomes.

1. Initial Assessment and Monitoring

Nurses play a key role in continuously assessing and monitoring the patient's neurological status, vital signs, and level of consciousness [5]. During the initial assessment, the nurse should collect important information about the patient's medical history, including any previous head injuries, current medications, and allergies [6].

Also, they should assess the severity of the trauma and document the Glasgow Coma Scale (GCS) score, which helps determine the level of consciousness and neurological functioning [7]. Additionally, vital signs such as blood pressure (BP), heart rate, and respiratory rate should be monitored continuously to detect any signs of deterioration. By closely monitoring these parameters, nurses can provide timely and effective care, optimize patient outcomes, and contribute to the overall management of tSAH patients [8].

1.1. Neurological Examinations

Neurological assessment is a fundamental aspect of caring for patients with tSAH. The nurse should closely monitor the patient's GCS, pupillary response, motor strength, and sensation [9]. Any changes in these parameters may indicate worsening intracranial pressure (ICP) or neurological deterioration [10]. Frequent examinations of cranial nerve function, including facial symmetry, eye movements, and gag reflex are important to identify any deficits that may require immediate intervention [11].

1.2. Monitoring Vital Signs

Monitoring vital signs and oxygenation status is crucial in managing tSAH. Close attention should be given to BP control to prevent rebleeding and/or exacerbation of cerebral edema [12]. However, hypotension should be avoided, as it can lead to cerebral hypoperfusion; hence, careful management of BP to maintain adequate cerebral perfusion pressure is necessary [13]. Also, oxygen saturation levels should be monitored continuously, and supplemental oxygen should be administered

as needed to ensure optimal tissue oxygenation [14].

1.3. Laboratory and Imaging Studies

Laboratory monitoring is also an integral part of nursing care for patients with tSAH. Serial blood tests, including complete blood count, coagulation profile, electrolytes, and arterial blood gases, are necessary to evaluate the patient's overall condition, detect any abnormalities, and guide appropriate interventions [15]. Serial imaging studies such as computed tomography (CT) scans and/or magnetic resonance imaging (MRI) may be performed to assess the extent of bleeding, identify underlying structural damage, or monitor treatment response [16, 17].

2. Interventions

2.1. Medication Administration

In some patients, medications are administered to manage symptoms, prevent complications, and promote recovery [18]. One of the primary medications commonly administered is analgesic [19]. tSAH can cause severe headaches and discomfort, and nurses must carefully assess and administer appropriate analgesics to alleviate the patient's pain [20]. In addition to pain management, nurses administer medications to prevent seizures [21]. Indeed, tSAH can increase the risk of seizures, and antiepileptic drugs are commonly prescribed to reduce this risk [22]. Nurses need sufficient knowledge regarding the specific antiepileptic medications used and their dosage requirements to ensure that patients receive the correct treatment [23]. Furthermore, medications to manage BP may be necessary for patients with tSAH. Indeed, elevated BP could worsen the condition and increase the risk of rebleeding [24]. Hence, nurses closely monitor the patient's BP and administer antihypertensive medications as prescribed by the healthcare provider to maintain stable BP levels [25]. Another important aspect of medication administration for patients with tSAH is preventing thromboembolic events [26]. These patients are at an increased risk of developing thrombosis, which can lead to stroke or other serious complications [27]. Nurses administer prophylactic anticoagulant medi-

cations such as low molecular weight heparin or fondaparinux to reduce the risk of thrombus formation while closely monitoring the patient for adverse effects [28]. Overall, medication administration for patients with tSAH requires nurses with the high level of skill as well as knowledge [29]. In other words, they should be familiar with proper drug dosages, potential side effects, drug interactions, and contraindications [30]. Additionally, nurses must adhere to medication administration protocols, double-check dosages, and ensure accurate documentation to maintain patient safety [31]. Hence, by effectively managing pain, preventing seizures, controlling BP, and minimizing thromboembolic events, nurses contribute significantly to the overall care and recovery of patients with tSAH.

2.2. ICP Monitoring

In the management of patients with tSAH, ICP monitoring plays a crucial role, and nurses play a significant part in its implementation and management [32]. ICP monitoring involves measuring the pressure within the skull to assess the status of brain tissue perfusion and detect any abnormalities that could lead to further complications [33]. Nurses are responsible for setting up and maintaining the ICP monitoring system. They ensure the equipment is correctly calibrated, properly placed, and connected to the patient's intraventricular or intraparenchymal catheter [34]. This process may require precision and attention to detail to avoid potential errors that could compromise the accuracy of ICP measurements. Regular monitoring and interpretation of ICP values are also essential nursing responsibilities [35]. Nurses frequently assess and document ICP readings, closely monitoring any fluctuations or abnormal trends [36]. By closely monitoring ICP levels, nurses could promptly identify and report any concerning changes to the healthcare team, enabling timely interventions to prevent secondary brain injuries.

Indeed, nurses are involved in implementing strategies to manage ICP within acceptable limits. They collaborate with other healthcare professionals, such as neurosurgeons and intensivists, to develop and execute individualized care plans [37]. These plans may include

elevating the head of the bed, maintaining adequate cerebral perfusion pressure, administering medications to control ICP, and managing ventilation parameters to optimize oxygenation and carbon dioxide levels [38, 39]. Furthermore, nurses provide ongoing education and support to patients and their families regarding the purpose and importance of ICP monitoring [40], which provides the rationale behind these interventions.

2.3. Fluid and Electrolyte Balance

Regarding various complications of tSAH, including cerebral edema and electrolyte imbalances, close monitoring and appropriate interventions to provide both fluid and electrolyte balance are essential for optimal patient outcomes [41]. Maintaining euvolemia prevents complications such as hypovolemia or hypervolemia [42]. Also, adequate hydration is necessary to optimize cerebral perfusion and prevent secondary brain injury. However, excessive fluid administration could contribute to cerebral edema and increased ICP [43]. Nurses should closely monitor vital signs, neurological status, urine output, and laboratory values to evaluate fluid levels accurately [44, 45]. They may collaborate with other healthcare team members, e.g., pharmacists, to determine appropriate fluid replacement strategies, which may include crystalloids or colloids [46].

Electrolyte balance is another critical aspect of patient care in tSAH. Electrolytes such as sodium, potassium, calcium, and magnesium play vital roles in cellular function, nerve conduction, and maintaining osmotic pressure [47]. tSAH can disrupt electrolyte balance due to various factors, including blood loss, diuretic therapy, or inadequate intake [48]. Hence, nurses must regularly monitor serum electrolyte levels and promptly intervene to correct any imbalances. For example, hyponatremia, or hypernatremia, can impact cerebral function and worsen patient outcomes [49]. So, nurses should administer electrolyte replacements or collaborate with the healthcare team to adjust medication dosages to restore and maintain electrolyte balance. In addition to monitoring fluid and electrolyte balance, nurses should consider factors such as renal function and urine output [50]. Maintaining

adequate renal perfusion is crucial to prevent acute kidney injury, which can further complicate the patient's condition [51]. By closely monitoring urine output and assessing renal function through laboratory tests, nurses can identify early signs of impaired renal function and implement appropriate interventions, including adjusting fluid administration, using diuretic medications, or collaborating with the healthcare team to address underlying causes [52].

2.4. Preventing Infections

By following specific guidelines and employing evidence-based practices, nurses can effectively reduce the incidence of infections in patients with tSAH [53].

Firstly, meticulous hand hygiene is necessary for preventing infections. Nurses should adhere to strict handwashing protocols, utilizing soap and water or alcohol-based sanitizers before and after any direct contact with patients [54]. This practice significantly reduces the transmission of pathogens and protects both the patient and the healthcare provider from potential infections [55].

Another vital strategy is to maintain a clean and sterile environment. Regular cleaning and disinfection of patient care areas, such as bedside tables, equipment, and surfaces, are essential [56]. Nurses should ensure that all equipment used for invasive procedures, such as catheters or drains, are appropriately sterilized and handled using aseptic techniques [57].

Also, implementing proper wound care techniques is crucial in preventing surgical site infections in patients with tSAH [58]. Indeed, nurses should meticulously assess and monitor the integrity of surgical incisions or any open wounds. Regular dressing changes using sterile techniques and appropriate antimicrobial agents help reduce the risk of infection [59, 60]. Additionally, closely monitoring for signs of infection, such as redness, swelling, or drainage, allows for early detection and intervention [57]. In addition, effective management of invasive devices, including urinary catheters and central venous lines, is also vital in infection prevention [61]. Nurses should follow strict protocols for their insertion, maintenance, and removal [61, 62]. Regular

monitoring for any signs of infection or complications related to these devices is crucial [61]. Indeed, early identification and prompt intervention can help prevent the spread of infection and subsequent complications [63]. Education and communication are important aspects of infection prevention. Nurses should educate patients and their families about the importance of infection control measures, including hand hygiene, wound care, and recognition of signs and symptoms of infection [64, 65]. Additionally, clear communication among the healthcare team regarding isolation precautions, patient status, and any specific infection risks is critical to ensure a cohesive approach [66].

2.5. Emotional Support

Managing patients with tSAH requires a comprehensive approach that addresses both the physical and emotional well-being of the individual. Emotional support strategies play a key role in helping patients cope with the psychological impact of their condition [67]. Remember, each patient's experience with tSAH is unique, and their emotional needs may differ. Tailoring emotional support strategies to individual preferences and circumstances is essential for effective management [68]. Collaborating with a multidisciplinary team that includes healthcare professionals, psychologists, and social workers can provide comprehensive care and support for patients with tSAH [67].

2.5.1. Provide Clear and Honest Communication

Establishing open and transparent communication with patients is essential. It is important to explain the diagnosis, treatment options, and potential outcomes in a clear and compassionate manner [68]. Offering reassurance and answering any questions and/or concerns could help reduce anxiety and foster trust [69].

2.5.2. Empathetic Listening and Validation

Patients with tSAH often experience various emotions, such as fear, confusion, frustration, and grief [70, 71]. Active listening and validating their feelings can make them feel understood and supported. Encouraging them to express their emotions and offering empathy

can help them process their trauma and adjust to their new reality [72].

2.5.3. Psychosocial Interventions

Engaging patients in psychosocial interventions, e.g., counseling, support groups, and therapy sessions, can be immensely beneficial [73]. These interventions provide a safe space for patients to share their experiences, learn coping strategies, and receive guidance from professionals who specialize in trauma-related care [74].

2.5.4. Education and Information

Providing educational resources about tSAH, its management, and recovery can encourage patients to participate actively in their care [75]. Knowledge about their condition can reduce anxiety, promote self-care, and enhance their understanding of the healing process [76, 77].

2.5.5. Encourage Social Support

Building a network of social support is crucial for patients with tSAH. Social support can provide emotional stability, practical assistance, and a sense of belonging, positively impacting the patient's overall well-being [78, 79].

2.5.6. Addressing Post-Traumatic Stress

Some patients may develop post-traumatic stress disorder (PTSD) following tSAH [80]. Recognizing the signs and symptoms of PTSD, such as flashbacks, nightmares, and avoidance behaviors, is essential [81]. Referring patients to mental health professionals who are specialized in trauma can help them receive appropriate treatment and support.

2.5.7. Holistic Approaches

Incorporating holistic approaches, such as relaxation techniques, mindfulness exercises, and complementary therapies, such as art and/or music therapy, can contribute to the emotional well-being of patients [82, 83]. These practices promote stress reduction, emotional regulation, and overall psychological resilience [83].

3. Rehabilitation Assistance

Early mobilization is a key strategy in reha-

bilitation assistance for tSAH patients [84]. Nurses work closely with physical therapists to ensure that patients start engaging in mobility exercises as soon as their medical condition allows [85]. This may involve helping the patient sit up, stand, or take their first steps. Early mobilization not only helps prevent complications such as muscle atrophy and joint stiffness but also promotes neuroplasticity and improves overall functional outcomes [86].

Communication assistance is another vital aspect of rehabilitation for tSAH patients. Many individuals with tSAH experience communication difficulties due to damage to the brain's language centers [87]. Nurses can collaborate with speech-language pathologists to implement strategies such as augmentative and alternative communication techniques, including using picture boards or electronic devices to facilitate effective communication [88]. Additionally, nurses can provide emotional support and patience during the patient's rehabilitation process, as frustration and anxiety may arise from the challenges of verbal expression [89].

Cognitive rehabilitation is also an essential component of managing tSAH patients. Nurses, alongside occupational therapists and neuropsychologists, can develop individualized cognitive training programs to improve attention, memory, problem-solving skills, and other cognitive functions [90, 91]. These programs may include puzzles, memory games, and computer-based exercises [92]. Nurses play a critical role in guiding and supporting patients throughout these sessions, monitoring their progress, and providing feedback to optimize outcomes. Also, psychosocial support is equally important in the rehabilitation of tSAH patients [93]. Nurses can assess and address the patient's emotional well-being, providing counseling and education to the patient and their families regarding the psychological consequences of tSAH. Also, they can facilitate support groups or connect patients with community resources to promote a sense of belonging and reduce feelings of isolation [94]. Additionally, nurses can collaborate with social workers to help address any financial or logistical challenges impeding the patient's rehabilitation progress [95].

4. Health Education and Discharge Planning

Health education begins immediately upon admission and continues throughout the patient's hospital stay [96]. Nurses aim to provide comprehensive information about tSAH, including its causes, risk factors, symptoms, and potential complications [97]. They explain the diagnostic tests, such as CT scans and cerebral angiography, and help patients understand their results. By promoting health literacy, nurses encourage patients to actively participate in their care and make informed decisions [98].

Furthermore, nurses educate patients about the treatment options available for tSAH. This may include surgical interventions, as well as medical management strategies to control BP, prevent vasospasm, and manage pain [99, 100]. They explain the purpose, benefits, and potential risks or side effects associated with each intervention, ensuring patients clearly understand their treatment plan.

Discharge planning is an integral part of managing patients with tSAH, as it involves coordinating services and resources to support the transition from the hospital to home or another healthcare setting [101]. Nurses collaborate with interdisciplinary teams, including physicians, physical therapists, occupational therapists, and social workers, to develop a comprehensive discharge plan tailored to the patient's individual needs [102]. During discharge planning, nurses assess the

patient's functional abilities, cognitive status, and psychosocial factors that may impact their recovery. Also, they provide guidance on medication management, wound care, activity restrictions, and lifestyle modifications to prevent recurrent bleeding or complications [103, 104].

In addition to physical care, nurses emphasize the importance of emotional support and coping strategies for patients and their families. They provide resources for counseling services or support groups to help individuals navigate the emotional impact of tSAH and its potential long-term effects [105].

Conclusion

Optimal nursing care plays a vital role in the management of patients with tSAH. Nurses contribute significantly to achieving positive patient outcomes through early recognition, comprehensive assessment, vigilant monitoring, implementation of therapeutic interventions, psychosocial support, patient education, and collaborative care. Ongoing research and evidence-based practice guidelines continue to shape nursing care strategies, leading to further improvements in managing patients with tSAH and their overall quality of life.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Witiw CD, Byrne JP, Nassiri F, Badhiwala JH, Nathens AB, Da Costa LB. Isolated traumatic subarachnoid hemorrhage: an evaluation of critical care unit admission practices and outcomes from a North American perspective. *Critical care medicine*. 2018 Mar 1;46(3):430-6.
2. Griswold DP, Fernandez L, Rubiano AM. Traumatic subarachnoid hemorrhage: a scoping review. *Journal of neurotrauma*. 2022 Jan 1;39(1-2):35-48.
3. Modi NJ, Agrawal M, Sinha VD. Post-traumatic subarachnoid hemorrhage: A review. *Neurology India*. 2016 Feb 1;64(7):8.
4. Tharayil AM, Chughtai TS, Younis B, Alyafei A, Verma V. Traumatic Subarachnoid Hemorrhage. *Management of Subarachnoid Hemorrhage*. 2022:179-88.
5. McNett MM, Gianakis A. Nursing interventions for critically ill traumatic brain injury patients. *Journal of Neuroscience Nursing*. 2010 Apr 1;42(2):71-7.
6. Ortega-Perez S, Shoyombo I, Aiyagari V, Atem F, Hill M, Stutzman SE, Olson DM. Pupillary light reflex variability as a predictor of clinical outcomes in subarachnoid hemorrhage. *Journal of Neuroscience Nursing*. 2019 Aug 1;51(4):171-5.
7. Ali AF, Sliman AM, Elnosary AM. Effect of Implementing Initial Neuroprotective Nursing Care on Outcomes of Traumatic Brain Injury Patients. *International Egyptian Journal of Nursing Sciences and Research*. 2023 Jul 29;4(1):370-84.

8. Fitzgerald E, Hammond N, Tian DH, Bradford C, Flower O, Harbor K, Johnson P, Lee R, Parkinson J, Tracey A, Delaney A. Functional outcomes at 12 months for patients with traumatic brain injury, intracerebral haemorrhage and subarachnoid haemorrhage treated in an Australian neurocritical care unit: A prospective cohort study. *Australian Critical Care*. 2020 Nov 1;33(6):497-503.
9. Evans V. Caring for traumatic brain injury patients: Australian nursing perspectives. *Critical Care Nursing Clinics*. 2021 Mar 1;33(1):21-36.
10. Özçelik EE, Celik S. Comparison of the Turkish versions of the Glasgow Coma Scale and four score used by intensive care nurses for neurological evaluation. *Journal of Clinical Nursing*. 2022 May;31(9-10):1397-406.
11. Čustović M. Monitoring in the Treatment of Patients with Aneurysmal and Nonaneurysmal Subarachnoid Haemorrhage in Terms of Nursing Care and Therapy. *Croatian Nursing Journal*. 2019 Jun 28;3(1):5-23.
12. Gamble M, Luggya TS, Mabweijano J, Nabulime J, Mowafi H. Impact of nursing education and a monitoring tool on outcomes in traumatic brain injury. *African journal of emergency medicine*. 2020 Dec 1;10(4):181-7.
13. Brice AE, Brice RG, Wallace SE. Recovery From a Subarachnoid Hemorrhage: Days 1 Through 22. *Communication Disorders Quarterly*. 2016 Nov;38(1):46-51.
14. Su Z, Guo W, Luo Y, Wang Y, Du Y. Observation on the Nursing Effect of the Whole Process in Patients with Severe Intracranial Hemorrhage. *Computational and Mathematical Methods in Medicine*. 2022 ;2022:1546019.
15. Butler JA. Treatment of subarachnoid haemorrhage complicated by hyponatraemia. *Australasian Journal of Neuroscience*. 2017 Oct 1;27(2):1-5.
16. Patel SK, Gozal YM, Krueger BM, Bayley JC, Moody S, Andaluz N, Falcone Jr RA, Bierbrauer KS. Routine surveillance imaging following mild traumatic brain injury with intracranial hemorrhage may not be necessary. *Journal of Pediatric Surgery*. 2018 Oct 1;53(10):2048-54.
17. Roguski M, Morel B, Sweeney M, Talan J, Rideout L, Riesenburger RI, Madan N, Hwang S. Magnetic resonance imaging as an alternative to computed tomography in select patients with traumatic brain injury: a retrospective comparison. *Journal of Neurosurgery: Pediatrics*. 2015 May 1;15(5):529-34.
18. Marcolini E, Hine J. Approach to the diagnosis and management of subarachnoid hemorrhage. *Western Journal of Emergency Medicine*. 2019 Mar;20(2):203.
19. Abraham MK, Chang WT. Subarachnoid hemorrhage. *Emergency Medicine Clinics*. 2016 Nov 1;34(4):901-16.
20. Jaffa MN, Podell JE, Foroutan A, Motta M, Chang WT, Cherian J, Pergakis MB, Parikh GY, Simard JM, Armahizer MJ, Badjatia N. Steroids Provide Temporary Improvement of Refractory Pain Following Subarachnoid Hemorrhage. *The Neurohospitalist*. 2023 Jul;13(3):236-42.
21. Boling W, Kore L. Subarachnoid hemorrhage-related epilepsy. *Subarachnoid Hemorrhage: Neurological Care and Protection*. 2020:21-5.
22. Feng R, Mascitelli J, Chartrain AG, Margetis K, Mocco J. Anti-epileptic drug (AED) use in subarachnoid hemorrhage (SAH) and intracranial hemorrhage (ICH). *Current Pharmaceutical Design*. 2017 Nov 1;23(42):6446-53.
23. Alkhachroum A, Megjhani M, Terilli K, Rubinos C, Ford J, Wallace BK, Roh DJ, Agarwal S, Connolly ES, Boehme AK, Claassen J. Hyperemia in subarachnoid hemorrhage patients is associated with an increased risk of seizures. *Journal of Cerebral Blood Flow & Metabolism*. 2020 Jun;40(6):1290-9.
24. Suarez JI. Diagnosis and management of subarachnoid hemorrhage. *CONTINUUM: Lifelong Learning in Neurology*. 2015 Oct 1;21(5):1263-87.
25. Carcel C, Sato S, Anderson CS. Blood pressure management in intracranial hemorrhage: current challenges and opportunities. *Current treatment options in cardiovascular medicine*. 2016 Apr;18:1-1.
26. Hantsche A, Wilhelmy F, Kasper J, Wende T, Hamerla G, Rasche S, Meixensberger J, Lindner D. Early prophylactic anticoagulation after subarachnoid hemorrhage decreases systemic ischemia and improves outcome. *Clinical Neurology and Neurosurgery*. 2021 Aug 1;207:106809.
27. Frisoli FA, Shinseki M, Nwabuobi L, Zeng XL, Agrados M, Kanter C, Frangos SG, Huang PP. Early venous thromboembolism chemoprophylaxis after traumatic intracranial hemorrhage. *Neurosurgery*. 2017 Dec 1;81(6):1016-20.
28. Miao W, Zhao K, Deng W, Teng J. Coagulation factor hyperfunction after subarachnoid hemorrhage induces deep venous thrombosis. *World Neurosurgery*. 2018 Feb 1;110:e46-52.
29. Mohamed Abdulmniem W, Mustafa Ragheb

- M, Saeid Mohamed S, Muhammed Hamed S. Effect of an Educational Program on Nurses Performance regarding Care for Patients with Subarachnoid Hemorrhage. *Journal of Nursing Science Benha University*. 2023 Jan 1;4(1):83-94.
30. Jin L, Sun L. Application of patient-centered care using guidelines of the Joint Commission on Accreditation of Healthcare Organizations in patients with acute subarachnoid hemorrhage. *American Journal of Translational Research*. 2021;13(4):2915.
31. Hutri I, Kotila J, Turunen H. Registered nurses' competence for subarachnoid haemorrhage patients in intensive care units. *British Journal of Neuroscience Nursing*. 2022 Oct 1;18(Sup5):S12-6.
32. Zoerle T, Lombardo A, Colombo A, Longhi L, Zanier ER, Rampini P, Stocchetti N. Intracranial pressure after subarachnoid hemorrhage. *Critical care medicine*. 2015 Jan 1;43(1):168-76.
33. Kramer AH. Critical ICP in subarachnoid hemorrhage: how high and how long?. *Neurocritical Care*. 2021 Jun;34(3):714-6.
34. Fiore M, Bogossian E, Creteur J, Oddo M, Taccone FS. Role of brain tissue oxygenation (PbtO₂) in the management of subarachnoid haemorrhage: a scoping review protocol. *BMJ open*. 2020 Sep 1;10(9):e035521.
35. Lee S, Kim YO, Baek JS, Ryu JA. The prognostic value of optic nerve sheath diameter in patients with subarachnoid hemorrhage. *Critical Care*. 2019 Dec;23(1):1-9.
36. Herzer G, Illievich U, Voelckel W, Trimmel H. Current practice in neurocritical care of patients with subarachnoid haemorrhage and severe traumatic brain injury. *Wiener klinische Wochenschrift*. 2016 ;128:649-657.
37. de Oliveira Manoel AL, Goffi A, Marotta TR, Schweizer TA, Abrahamson S, Macdonald RL. The critical care management of poor-grade subarachnoid haemorrhage. *Critical care*. 2016 Dec;20(1):1-9.
38. Kienzler JC, Bäbler S, Zakelis R, Remonda E, Ragauskas A, Fandino J. Validation of noninvasive absolute intracranial pressure measurements in traumatic brain injury and intracranial hemorrhage: preliminary results. *Journal of Neurological Surgery Part A: Central European Neurosurgery*. 2015 Sep;76(S 01):P065.
39. Lv Y, Wang D, Lei J, Tan G. Clinical observation of the time course of raised intracranial pressure after subarachnoid hemorrhage. *Neurological Sciences*. 2015 Jul;36:1203-10.
40. Ravishankar N, Nuoman R, Amuluru K, El-Ghanem M, Thulasi V, Dangayach NS, Lee K, Al-Mufti F. Management strategies for intracranial pressure crises in subarachnoid hemorrhage. *Journal of intensive care medicine*. 2020 Mar;35(3):211-8.
41. Suman S, Kumar N, Singh Y, Kumar V, Yadav G, Gupta BK, Pandey AR, Pandey S. Evaluation of serum electrolytes in traumatic brain injury patients: Prospective randomized observational study. *J Anesth Crit Care Open Access*. 2016 Jul 28;5(3):00184.
42. Pin-On P, Saringkarinkul A, Punjasawadwong Y, Kacha S, Wilairat D. Serum electrolyte imbalance and prognostic factors of postoperative death in adult traumatic brain injury patients: A prospective cohort study. *Medicine*. 2018 ;97(45):e13081.
43. Schur S, Panagiotoglou D, Di Maio S, Buckeridge D. Comorbidities and medical complications in hospitalized subarachnoid hemorrhage patients. *Canadian Journal of Neurological Sciences*. 2022 Jul;49(4):569-78.
44. Chen I, Mitchell P. Serum potassium and sodium levels after subarachnoid haemorrhage. *British Journal of Neurosurgery*. 2016 Sep 2;30(5):554-9.
45. Connor Jr DE, Chaitanya GV, Chittiboina P, McCarthy P, Scott LK, Schrott L, Minagar A, Nanda A, Alexander JS. Variations in the cerebrospinal fluid proteome following traumatic brain injury and subarachnoid hemorrhage. *Pathophysiology*. 2017 Sep 1;24(3):169-83.
46. Neligan PJ. Fluid and electrolyte balance. *Anaesthesia & Intensive Care Medicine*. 2021 Mar 1;22(3):169-73.
47. Nwafor DC, Kirby BD, Ralston JD, Colantonio MA, Ibekwe E, Lucke-Wold B. Neurocognitive Sequelae and Rehabilitation after Subarachnoid Hemorrhage: Optimizing Outcomes. *Journal of vascular diseases*. 2023 Apr 1;2(2):197-211.
48. Yamashita RH, Yamaki VN, Rabelo NN, Welling LC, Figueiredo EG. Acid-Base and Electrolyte Disorders in Neurocritical Care. *Neurocritical Care for Neurosurgeons: Principles and Applications*. 2021:373-90.
49. Mezzini G, Marasco S, Bertuccio A, Savioli G, Piccolella F, Racca F, Barbanera A, Vitali M. Hyponatremia Related to Neurocritical Care: Focus on Diagnosis and Therapy: A Systematic Review. *Reviews on Recent Clinical Trials*. 2023 Feb 1;18(1):19-27.
50. Sadan O, Singbartl K, Kandiah PA, Martin KS, Samuels OB. Hyperchloremia is associated with acute kidney injury in patients with subarachnoid hemorrhage. *Critical care*

- medicine. 2017 Aug 1;45(8):1382-8.
51. Farrokh S, Cho SM, Suarez JI. Fluids and hyperosmolar agents in neurocritical care: an update. *Current opinion in critical care*. 2019 Apr 1;25(2):105-9.
 52. Shah K, Turgeon RD, Gooderham PA, Ensom MH. Prevention and treatment of hyponatremia in patients with subarachnoid hemorrhage: a systematic review. *World Neurosurgery*. 2018 Jan 1;109:222-9.
 53. Bogossian EG, Attanasio L, Creteur J, Grimaldi D, Schuind S, Taccone FS. The impact of extracerebral infection after subarachnoid hemorrhage: A single-center cohort study. *World Neurosurgery*. 2020 Dec 1;144:e883-97.
 54. Rhim HY, Won SY, Kashefiolasl S, Brawanski N, Hattingen E, Berkefeld J, Seifert V, Konczalla J. Multidrug-resistant organisms (MDROs) in patients with subarachnoid hemorrhage (SAH). *Scientific Reports*. 2021 Apr 15;11(1):8309.
 55. Kaplan A, Kaleem S, Huynh M. Quality Improvement in the Management of Subarachnoid Hemorrhage: Current State and Future Directions. *Current Pain and Headache Reports*. 2023 Mar;27(3):27-38.
 56. Kofler M, Beer R, Marinoni S, Schiefecker AJ, Gaasch M, Rass V, Lindner A, Lanosi BA, Rhomberg P, Pfaußler B, Thomé C. Early supplemental parenteral nutrition for the achievement of nutritional goals in subarachnoid hemorrhage patients: An observational cohort study. *Plos one*. 2022 Mar 18;17(3):e0265729.
 57. Roa'a WJ, Lahlouh AB, Alshogran OY, Aldabbour BA, Balusha AA. Nosocomial infections among patients with intracranial hemorrhage: a retrospective data analysis of predictors and outcomes. *Clinical Neurology and Neurosurgery*. 2019 Jul 1;182:158-66.
 58. Al-Mufti F, Amuluru K, Changa A, Lander M, Patel N, Wajswol E, Al-Marsoum S, Alzubaidi B, Singh IP, Nuoman R, Gandhi C. Traumatic brain injury and intracranial hemorrhage-induced cerebral vasospasm: A systematic review. *Neurosurgical focus*. 2017 Nov 1;43(5):E14.
 59. Andrews PJ, Verma V, Healy M, Lavinio A, Curtis C, Reddy U, Andrzejowski J, Foulkes A, Canestrini S. Targeted temperature management in patients with intracerebral haemorrhage, subarachnoid haemorrhage, or acute ischaemic stroke: consensus recommendations. *British Journal of Anaesthesia*. 2018 Oct 1;121(4):768-75.
 60. Lissak IA, Locascio JJ, Zafar SF, Schleicher RL, Patel AB, Leslie-Mazwi T, Stapleton CJ, Koch MJ, Kim JA, Anderson K, Rosand J. Electroencephalography, hospital complications, and longitudinal outcomes after subarachnoid hemorrhage. *Neurocritical Care*. 2021 Jan 22:1-2.
 61. Rabinstein AA, Sandhu K. Non-infectious fever in the neurological intensive care unit: incidence, causes and predictors. *Journal of Neurology, Neurosurgery & Psychiatry*. 2007 Nov 1;78(11):1278-80.
 62. van Gijn J, Rinkel GJ. Subarachnoid haemorrhage: diagnosis, causes and management. *Brain*. 2001 Feb 1;124(2):249-78.
 63. Friedrich V, Flores R, Muller A, Bi W, Peerschke EI, Sehba FA. Reduction of neutrophil activity decreases early microvascular injury after subarachnoid haemorrhage. *Journal of neuroinflammation*. 2011 Dec;8:1-2.
 64. Lee S, Kim YO, Ryu JA. Clinical usefulness of early serial measurements of C-reactive protein as outcome predictors in patients with subarachnoid hemorrhage. *BMC neurology*. 2020 Dec;20:1-0.
 65. Wu Y, Liu Y, Zhou C, Wu Y, Sun J, Gao X, Huang Y. Biological effects and mechanisms of caspases in early brain injury after subarachnoid hemorrhage. *Oxidative Medicine and Cellular Longevity*. 2022 ;2022:3345637.
 66. Chou SH. Subarachnoid hemorrhage. *CONTINUUM: Lifelong Learning in Neurology*. 2021 Oct 1;27(5):1201-45.
 67. Taufique Z, May T, Meyers E, Falo C, Mayer SA, Agarwal S, Park S, Connolly ES, Claassen J, Schmidt JM. Predictors of poor quality of life 1 year after subarachnoid hemorrhage. *Neurosurgery*. 2016 Feb 1;78(2):256-64.
 68. Bartlett M, Bulters D, Hou R. Psychological distress after subarachnoid haemorrhage: A systematic review and meta-analysis. *Journal of Psychosomatic Research*. 2021 Sep 1;148:110559.
 69. Tsyben A, Guilfoyle MR, Laing RJ, Timofeev I, Anwar F, Trivedi RA, Kirolos RW, Turner C, Allanson J, Mee H, Outtrim JG. Comparison of health-related quality of life in patients with traumatic brain injury, subarachnoid haemorrhage and cervical spine disease. *British journal of neurosurgery*. 2022 Dec 8:1-7.
 70. Rueckriegel SM, Baron M, Domschke K, Neuderth S, Kunze E, Kessler AF, Nickl R, Westermaier T, Ernestus RI. Trauma- and distress-associated mental illness symptoms in close relatives of patients with severe traumatic brain injury and high-grade subarachnoid hemorrhage. *Acta*

- neurochirurgica. 2015 Aug;157:1329-36.
71. Edlow BL, Samuels O. Emergency neurological life support: subarachnoid hemorrhage. *Neurocritical Care*. 2017 Sep;27:116-23.
 72. Ditty BJ, Omar NB, Foreman PM, Patel DM, Pritchard PR, Okor MO. The nonsurgical nature of patients with subarachnoid or intraparenchymal hemorrhage associated with mild traumatic brain injury. *Journal of neurosurgery*. 2015 Sep 1;123(3):649-53.
 73. Tang WK, Wang L, Wong GK, Ungvari GS, Yasuno F, Tsoi KK, Kim JS. Depression after subarachnoid hemorrhage: a systematic review. *Journal of stroke*. 2020 Jan;22(1):11.
 74. Tang WK, Wang L, Tsoi KK, Barrash J, Kim JS. Personality changes after subarachnoid hemorrhage: A systematic review and meta-analysis. *Journal of psychosomatic research*. 2022 May 1;156:110762.
 75. Zhang H, Zhang M, Zhao F. Nursing Method of Patients with Severe Traumatic Brain Injury and Fracture in the Ambulance. *BioMed Research International*. 2022;2022:2652916.
 76. Byun E, McCurry SM, Kim B, Kwon S, Thompson HJ. Sleep disturbance and self-management in adults with subarachnoid hemorrhage: a qualitative study. *Clinical Nursing Research*. 2022 May;31(4):632-8.
 77. Tang WK, Wang L, Tsoi KK, Kim JM, Lee SJ, Kim JS. Anxiety after subarachnoid hemorrhage: A systematic review and meta-analysis: Anxiety in subarachnoid hemorrhage. *Journal of Affective Disorders Reports*. 2021 Jan 1;3:100060.
 78. McIntosh AP, Thomas A. Health-related quality-of-life outcomes: comparing patients with aneurysmal and nonaneurysmal subarachnoid hemorrhage. *Journal of Neuroscience Nursing*. 2015 Oct 1;47(5):E2-11.
 79. Brice RG, Brice A. Recovery From a Subarachnoid Hemorrhage: Patient and Spouse Perspectives. *Communication Disorders Quarterly*. 2017 Feb;38(2):119-28.
 80. Tang WK, Wang L, Tsoi KK, Kim JS. Post-Traumatic Stress Disorder after Subarachnoid Hemorrhage: A Systematic Review. *Neurology India*. 2023 Jan 1;71(1):9.
 81. Garton AL, Gupta VP, Pucci JU, Couch CK, Connolly Jr ES. Incidence and predictors of post-traumatic stress symptoms in a cohort of patients with intracerebral hemorrhage. *Clinical Neurology and Neurosurgery*. 2020 Mar 1;190:105657.
 82. Hutri I, Kotila J, Turunen H. Registered nurses' competence for subarachnoid haemorrhage patients in intensive care units. *British Journal of Neuroscience Nursing*. 2022 Oct 1;18(Sup5):S12-6.
 83. Compton ES, Smallheer BA, Thomason NR, Norris MS, Nordness MF, Smith MD, Patel MB. Minimal-risk traumatic brain injury management without neurosurgical consultation. *Journal of Neurocritical Care*. 2020 Dec 14;13(2):80-5.
 84. Sorek G, Shaklai S, Gagnon I, Schneider K, Chevnard M, Stern N, Fadida Y, Kalderon L, Katz-Leurer M. Impact of Subarachnoid Hemorrhage on the Cardiac Autonomic Function During Rehabilitation in Children After Severe Traumatic Brain Injury. *Neurotrauma Reports*. 2023 Jul 1;4(1):458-62.
 85. Cinotti R, Putegnati JB, Lakhali K, Desal H, Chenet A, Buffenoir K, Frasca D, Allaouchiche B, Asehnoune K, Rozec B. Evolution of neurological recovery during the first year after subarachnoid haemorrhage in a French university centre. *Anaesthesia Critical Care & Pain Medicine*. 2019 Jun 1;38(3):251-7.
 86. Konczalla J, Schmitz J, Kashefiolasi S, Senft C, Platz J, Seifert V. Non-aneurysmal non-perimesencephalic subarachnoid hemorrhage: effect of rehabilitation at short-term and in a prospective study of long-term follow-up. *Topics in Stroke Rehabilitation*. 2016 Jun 6;23(4):261-8.
 87. Lindner A, Brunelli L, Rass V, Ianosi BA, Gaasch M, Kofler M, Limmert V, Schiefecker AJ, Pfausler B, Beer R, Pucks-Faes E. Long-term clinical trajectory of patients with subarachnoid hemorrhage: linking acute care and neurorehabilitation. *Neurocritical Care*. 2023 Feb;38(1):138-48.
 88. Reznik ME, Schmidt JM, Mahta A, Agarwal S, Roh DJ, Park S, Frey HP, Claassen J. Agitation after subarachnoid hemorrhage: a frequent omen of hospital complications associated with worse outcomes. *Neurocritical care*. 2017 Jun;26:428-35.
 89. Tölli A, Höybye C, Bellander BM, Johansson F, Borg J. The effect of time on cognitive impairments after non-traumatic subarachnoid haemorrhage and after traumatic brain injury. *Brain injury*. 2018 Oct 15;32(12):1465-76.
 90. Boerboom W, Heijnenbroek-Kal MH, Van Kooten F, Khajeh L, Ribbers GM. Unmet needs, community integration and employment status four years after subarachnoid haemorrhage. *Journal of Rehabilitation Medicine*. 2016 May 20;48(6):529-34.
 91. Persson HC, Törnblom K, Sunnerhagen KS, Törnblom M. Consequences and coping strategies six years after a subarachnoid hemorrhage—A qualitative study. *PLoS One*. 2017 Aug 30;12(8):e0181006.

92. Reznik ME, Mahta A, Schmidt JM, Frey HP, Park S, Roh DJ, Agarwal S, Claassen J. Duration of agitation, fluctuations of consciousness, and associations with outcome in patients with subarachnoid hemorrhage. *Neurocritical care*. 2018 Aug;29:33-9.
93. Zeiler FA, Lo BW, Akoth E, Silvaggio J, Kaufmann AM, Teitelbaum J, West M. Predicting outcome in subarachnoid hemorrhage (SAH) utilizing the Full Outline of UnResponsiveness (FOUR) score. *Neurocritical care*. 2017 Dec;27:381-91.
94. Sonesson B, Kronvall E, Säveland H, Brandt L, Nilsson OG. Long-term reintegration and quality of life in patients with subarachnoid hemorrhage and a good neurological outcome: findings after more than 20 years. *Journal of neurosurgery*. 2017 Apr 28;128(3):785-92.
95. Eapen BC, Allred DB, O'Rourke J, Cifu DX. Rehabilitation of moderate-to-severe traumatic brain injury. In *Seminars in neurology*. 2015; 35(1): 1-13.
96. Collins CI, Hasan TF, Mooney LH, Talbot JL, Fouraker AL, Nelson KF, Ohanian M, Bonnett SL, Tawh RG, Nordan LM, Hodge DO. Subarachnoid hemorrhage "fast track": a health economics and health care redesign approach for early selected hospital discharge. *Mayo Clinic Proceedings: Innovations, Quality & Outcomes*. 2020 Jun 1;4(3):238-48.
97. Yousefzadeh-Chabok S, Kapourchali FR, Ramezani S. Determinants of long-term health-related quality of life in adult patients with mild traumatic brain injury. *European journal of trauma and emergency surgery*. 2021 Jun;47:839-46.
98. Weimer JM, Nowacki AS, Frontera JA. Withdrawal of life-sustaining therapy in patients with intracranial hemorrhage: self-fulfilling prophecy or accurate prediction of outcome?. *Critical Care Medicine*. 2016 Jun 1;44(6):1161-72.
99. Byun E, McCurry SM, Opp M, Liu D, Becker KJ, Thompson HJ. Self-efficacy is associated with better sleep quality and sleep efficiency in adults with subarachnoid hemorrhage. *Journal of Clinical Neuroscience*. 2020 Mar 1;73:173-8.
100. Kurogi R, Kada A, Ogasawara K, Nishimura K, Kitazono T, Iwama T, Matsumaru Y, Sakai N, Shiokawa Y, Miyachi S, Kuroda S. National trends in the outcomes of subarachnoid haemorrhage and the prognostic influence of stroke centre capability in Japan: retrospective cohort study. *BMJ open*. 2023 Apr 1;13(4):e068642.
101. Koso M, Dizdarevic K. Attention and executive functions in microsurgically treated patients after subarachnoid hemorrhage. *Asian Journal of Neurosurgery*. 2015 Dec;10(04):260-5.
102. Macdonald RL. Subarachnoid Hemorrhage. *Handbook of Neuroemergency Clinical Trials*. 2018 Jan 1:23-45.
103. Joseph B, Obaid O, Dultz L, Black G, Campbell M, Berndtson AE, Costantini T, Kerwin A, Skarupa D, Burruss S, Delgado L. Validating the Brain Injury Guidelines: Results of an American Association for the Surgery of Trauma prospective multi-institutional trial. *Journal of trauma and acute care surgery*. 2022 Aug 1;93(2):157-65.
104. Paget LM, Boutonnet M, Moyer JD, Delhaye N, D'Aranda E, Beltzer N, Hamada SR. Trauma centre admissions for traumatic brain injury in France: One-year epidemiological analysis of prospectively collected data. *Anaesthesia Critical Care & Pain Medicine*. 2021 Feb 1;40(1):100804.
105. Ogden JA, Mee EW, Henning M. A prospective study of impairment of cognition and memory and recovery after subarachnoid hemorrhage. *Neurosurgery*. 1993 Oct 1;33(4):572-87.