

ORIGINAL
ARTICLE

Received 2024-08-06
Revised 2024-09-28
Accepted 2024-11-08

Investigating the Morphology of the Nasal Cavity for Nasal Reconstruction Using Cone Beam Computed Tomography Images

Aisan Ghaznavi¹, Sina Ilkhani², Asma Sodaii¹, Mohammad Jafari Heydarlou¹ ✉

¹ Department of Oral and Maxillofacial Radiology, School of Dentistry, Urmia University of Medical Sciences, Urmia, Iran

² Department of Oral and Maxillofacial Surgery, School of Dentistry, Urmia University of Medical Sciences, Urmia, Iran

Abstract

Background: Facial reconstruction is the procedure of rebuilding a face onto an anonymous skull to aid identification in forensic and archaeological cases. This study investigated the morphology of the nasal cavity for reconstruction by using cone beam computed tomography images. **Materials and Methods:** In this retrospective cross-sectional study, pre-treatment CBCT images of 220 adults were selected by random sampling from the records of orthodontic clinical data between January 2022 and November 2023. The three-dimensional parameters of the nasal soft structures and hard structures were measured. **Results:** Of 220 CBCT images, 198 cases (61.1% females and 38.9% males) were examined in the final analysis after meeting inclusion criteria. TStatistically significant sex differences were observed in nasal length (males: 50.79 ± 4.78 mm, females: 45.28 ± 4.18 mm; $P < 0.05$), nasal depth (males: 23.54 ± 2.43 mm, females: 26.05 ± 3.53 mm; $p < 0.05$), and nasal height (males: 49.81 ± 4.30 mm, females: 55.02 ± 4.49 mm; $P < 0.05$). The nasolabial angle was significantly higher in males ($98.21^\circ \pm 8.34^\circ$) compared to females ($89.71^\circ \pm 7.37^\circ$; $P < 0.05$). Conversely, the nasal tip angle was significantly higher in females ($77.18^\circ \pm 8.45^\circ$) than in males ($71.54^\circ \pm 8.20^\circ$; $P < 0.05$). A statistically significant difference was also observed in the nasal upward tip angle between males ($23.8 \pm 3.10^\circ$) and females ($20.45^\circ \pm 2.98^\circ$; $P < 0.05$). **Conclusion:** This study revealed significant sex-based variations in nasal parameters. Males exhibited greater nasal length, depth, and nasal tip angle compared to females. [GMJ.2024;13:e3522] DOI: [10.31661/gmj.v13iSP1.3522](https://doi.org/10.31661/gmj.v13iSP1.3522)

Keywords: Nasal Morphology; Cone Beam Computed Tomography; Nasal Reconstruction; Nasal Cavity; Facial Index

Introduction

The nasal cavity, a intricate air-filled chamber, exerts a critical influence on three key functions: respiration, olfaction (smell), and facial aesthetics and does a vital position within the facial skeleton [1]. Although relatively stable during adolescence [2], nasal

development concludes by age 16 in females and 18 in males [3]. The intricate shape of the nasal cavity, defined by various bone and cartilage structures, plays a role in airflow, drainage, and ultimately, how well we breathe [4]. Trauma, birth defects, and tumors can disrupt this form, causing problems with both function and appearance [5].

GMJ

Copyright© 2024, Galen Medical Journal.
This is an open-access article distributed
under the terms of the Creative Commons
Attribution 4.0 International License
(<http://creativecommons.org/licenses/by/4.0/>)
Email: gmj@salviapub.com



✉ Correspondence to:

Mohammad Jafari Heydarlou, Department of Oral and Maxillofacial Radiology, School of Dentistry, Urmia University of Medical Sciences, Urmia, Iran.
Telephone Number: +98 9372308482
Email Address: dr.jafarymo@yahoo.com

Nasal reconstruction surgery exists to restore normalcy in these situations, aiming to improve both breathing and facial aesthetics [6]. Facial reconstruction has been extensively employed for craniofacial recognition and identification, as evidenced by numerous investigations [7]. A substantial body of research has quantified the relationship between the skull's skeletal structure and the overlying facial soft tissues [8-10]. This methodology, adaptable to the available skeletal remains and their condition, can be a valuable tool in aiding case identification, particularly in large-scale disasters. Prior studies have primarily relied on cadaveric analyses to investigate the intricate link between facial hard and soft tissues [11, 12]. Notably, nasal morphology exhibits high variability and plays a significant role in facial recognition due to its unique characteristics [13].

Therefore, accurate prediction of the nasal tip has the potential to enhance the overall accuracy of facial reconstructions for identification purposes [14]. The prediction of nasal morphology from the skull has been a topic of considerable investigation, as evidenced by numerous reports [15, 16]. Evaluation of nasal morphology has historically relied on various techniques, including morphometry, photogrammetry, radiography, and more recently, three-dimensional (3D) imaging modalities [17, 18].

Traditional methods such as lateral cephalom-

etry and plain radiography provided limited two-dimensional (2D) views, hindering the accurate assessment of the inherently complex 3D structures of the nasal cavity. However, the emergence of cone beam computed tomography (CBCT) has revolutionized nasal cavity imaging by offering high-resolution, 3D reconstructions. This advancement allows for a more comprehensive and accurate evaluation of nasal morphology.

To our knowledge, few studies have focused on the 3D spatial relationships between the hard tissue and overlying soft tissue of the CBCT images for nose reconstruction in Iranian ethnicity. The present study aimed to investigate the morphology of the nasal cavity for reconstruction by using cone beam computed tomography images.

Materials and Methods

In this analytical cross-sectional study, by random sampling, pre-treatment CBCT images of 220 Iranian adults with ages more than 18 years were selected from the records of a radiography centers affiliated to Uremia University of Medical Sciences from January 2022 to November 2023.

Adherence to ethical and legal requirements was paramount, and informed consent was obtained from all participants. Ethical approval was given by the medical ethics committee of Urmia University of Medical Sciences, the




Figure 1. Reference anthropological hard tissue landmarks.

following reference number was IR.UMSU.REC.1402.216.

To ensure the internal validity of the study, a strict exclusion criterion was employed. Patients with pre-existing medical conditions potentially affecting nasal morphology were excluded, including those with bone diseases, a history of nasal trauma, congenital nasal anomalies, prior nasal and facial sinus surgeries, and relevant nasal or maxillary sinus pathologies. Additionally, patients with known syndromes were excluded.

This study employed vertical cephalic radiographs with subsequent 3D reconstructions for data collection. The axial sections ensured complete visualization of the maxilla and its pterygoid processes. Scanning parameters were adjusted to optimize image quality based on individual patient characteristics. Two independent evaluators, a radiologist and a student, assessed the images using Ramixis 5.1.1 software on a computer equipped with a high-resolution (1920 x 1080 pixels) 42-inch LG monitor. Evaluations were conducted in a controlled environment free from light pollution. To assess nasal morphology, a set of established anthropological bony landmarks was utilized [15].

1- Sella (S): The midpoint of the pituitary fossa (Figure-1)

2- Nasion (N): The most anteriorly projecting point on the nasofrontal suture in the sagittal plane (Figure-1)

3- Pogonion (Pog): The most prominent anterior aspect of the mandibular symphysis (Figure-1)

4- Menton (M): The inferiormost point of the mandibular symphysis (Figure-1)

5- Rhinion (Rh): The most inferior point where the right and left nasal bones meet, essentially marking the junction between the bony and cartilaginous portions of the nasal structure in the midline (Figure-2).

6- Nasal Cavity (NC): The most laterally positioned point on the bony outline of the nasal cavity (Figure-3)

7- Anterior Nasal Spine (ANS): The anterior-most tip of the bony projection arising from the maxilla, located at the inferior border of the anterior nasal opening (Figure-3)

To comprehensively assess nasal morphology, this study additionally employed a set of established anthropological soft tissue landmarks [19] (Figure-4). These landmarks included:

1- Soft Tissue Nasion (N'): The deepest indentation in the midline between the brow ridge (glabella) and the tip of the soft tissue nose (pronasale).

2- Pronasale (Prn): The most forward-projecting point of the soft tissue nose, identified as the intersection of a perpendicular line drawn from the Frankfort Horizontal Plane (FHP) and the nasal tip.

3- Subnasale (Sn): The deepest midline point where the columella (the central pillar be-




Figure 2. Reference anthropological hard tissue landmarks.

tween the nostrils) meets the upper lip.

4- Posterior Columella Point (PCm): The rearmost point on the lower border of the nose where it curves downward to meet the philtral ridge of the upper lip.

5- Columella Point (Cm): The most anterior aspect of the columella.

6- Labrale Superius (Ls): The point marking the boundary between the skin and the mucous membrane on the upper lip.

7- Ala (Al): The outermost point on the outline of each nostril (ala).

This study employed a set of established reference planes and anthropometric variables to quantify nasal morphology [20]. These included:

1- Frankfort Horizontal Plane (FHP): A reference plane constructed by connecting the highest point on the opening of each ear canal (external auditory meatus) with the lowest point on the inferior margin of the left eye socket (orbit).

2- Nasal Length: The linear distance measured between the soft tissue nasion (N') and the pronasale (Prn).

3- Nasal Depth: The perpendicular distance between the pronasale (Prn) and a line drawn from the soft tissue nasion (N') to the subnasale (Sn).

4- Nasal Width (Ala-Ala): The maximum horizontal distance between the left and right alar points (outermost points of the nostrils).

5- Nasal Height (N' - Sn): The linear distance measured between the soft tissue nasion (N')

and the subnasale (Sn).

6- Nasal Index (Ala-Ala/N-Sn): A dimensionless ratio calculated by multiplying the ratio of the maximum nasal width (ala-ala) to the maximum nasal height (N' - Sn) by 100. This index helps categorize nasal breadth relative to height.

7- Nasolabial Angle (NLA): The angle formed at the intersection of a tangent line drawn along the posterior columella point (PCm) and a line connecting the PCm to the labrale superius (Ls) (the border between the upper lip's skin and mucous membrane).

8- Nasal Upward Tip Angle (UNLA): The angle formed when the tangent line drawn along the posterior columella point (PCm) is extended anteriorly to intersect the Frankfort Horizontal Plane (FHP). This angle reflects the upward rotation of the nasal tip.

9- Nasal Tip Angle (NTP): The angle formed by the intersection of the axis of the nasal dorsum (bridge) and the tangent line drawn along the posterior columella point (PCm). This angle reflects the projection of the nasal tip.

Sample Size Calculation

The sample size in the present study was calculated based on the study by Gruszka *et al.* [21], in which the ratio (p) of the most common changes related to the anatomical variation related to nasal cells was 54.8 and the confidence interval coefficient 95% $z_{(1-\alpha/2)}=1.96 \approx 2$ and the accuracy (d) for estimating the point which is the minimum important




Figure 3. Reference anthropological hard tissue landmarks.

difference from the clinical aspect for the difference between the groups was estimated as 198 according to the following formula.

Statistical Analysis

Quantitative findings were reported in the form of mean and standard deviation, and qualitative findings, which were the gender distribution of the studied subjects, were reported in the form of percentages. An independent samples t-test was used to compare the estimates between the males and females. All statistical analyses were performed using SPSS 22.0. $P < 0.05$ was considered to be statistically significant.

Results

To assess intra-observer variability, a random selection of 20 CBCT images were traced twice by the researcher at a two-week interval. The Kappa coefficient was calculated, demonstrating acceptable agreement (values not shown).

Prior to commencing the main study, a pilot evaluation was conducted with one radiologist and one student examiner following their training.

The intraclass correlation coefficient was used to assess inter-observer agreement, yielding values of 0.85, 0.92, and 0.91 for nasal length, depth, and height measurements, respectively. These values indicate good to excellent reliability.

A total of 220 CBCT images were initially examined. Following application of inclusion

and exclusion criteria, 198 cases were included in the final analysis. The sample comprised 121 females (61.1%) and 77 males (38.9%). Statistically significant sex differences were observed in nasal length (males: 50.79 ± 4.78 mm, females: 45.28 ± 4.18 mm; $p < 0.05$), nasal depth (males: 23.54 ± 2.43 mm, females: 26.05 ± 3.53 mm; $p < 0.05$), and nasal height (males: 49.81 ± 4.30 mm, females: 55.02 ± 4.49 mm; $p < 0.05$). However, no significant sex differences were found in nasal width (males: 25.95 ± 1.80 mm, females: 24.16 ± 2.13 mm; $p = 0.432$) or nasal index (males: $65.67 \pm 7.61\%$, females: $64.88 \pm 7.15\%$; $p = 0.467$). The nasolabial angle was significantly higher in males ($98.21^\circ \pm 8.34^\circ$) compared to females ($89.71^\circ \pm 7.37^\circ$; $p < 0.05$). Conversely, the nasal tip angle was significantly higher in females ($77.18^\circ \pm 8.45^\circ$) than in males ($71.54^\circ \pm 8.20^\circ$; $p < 0.05$). A statistically significant difference was also observed in the nasal upward tip angle between males ($23.8^\circ \pm 3.10^\circ$) and females ($20.45^\circ \pm 2.98^\circ$; $P < 0.05$, Table-1).

Discussion

Facial reconstruction is a multifaceted discipline that integrates scientific methodologies with artistic expertise to recreate soft tissues on a skull [22]. This technique offers valuable applications in forensic medicine, enabling the generation of facial approximations for identification and diagnostic purposes [23]. Furthermore, it plays a crucial role in facial reconstruction and corrective surgeries [24].

Table 1. Descriptive Data and Gender Variation of Nasal Parameters

	Male (n = 77 (38.9%))		Female (n = 121 (61.1%))		P-value
	Mean	SD	Mean	SD	
Nasal length (mm)	50.79	4.78	45.28	4.18	<0.000
Nasal depth (mm)	23.54	2.43	26.05	3.53	<0.000
Nasal width (mm)	25.95	1.8	24.16	2.13	0.432
Nasal height (mm)	49.81	4.3	55.02	4.49	<0.000
Nasal Index (%)	65.67	7.61	64.88	7.15	0.467
Nasolabial angle (NLA)	98.21	8.34	89.71	7.37	0.051
Nasal upward tip angle (UNLA)	23.8	3.1	20.45	2.98	<0.000
Nasal tip angle (NTP)	78.18	8.45	71.54	8.2	<0.000

In light of this, the present study employed CBCT images to investigate nasal cavity morphology in the context of facial reconstruction.

Previous research has employed diverse methodologies and anatomical landmarks to evaluate nasal morphology and its relationship with surrounding hard and soft tissues, primarily using lateral cephalometric radiographs [17, 25, 15]. For instance, Aljabaa *et al.* investigated the influence of sex and ethnicity on nasal form and its association with other cranial structures, highlighting the importance of these factors when establishing normative values [17]. Additionally, researchers have adopted quantitative approaches to assess nasal morphology and its connections to other facial elements within distinct ethnic populations [15, 25].

However, critiques have emerged regarding the application of specific methods. Lapointe *et al.* argued for the use of Stephan's craniofacial norm instead of Gerasimov's method, suggesting potential inaccuracies in the latter [26]. Furthermore, studies focusing on specific Asian populations have contributed valuable data. Other related study provided detailed information on craniofacial soft tissue thickness and nasal profile characteristics within the Chinese Xi'an Han population [27].

Advancements in dental technology have introduced innovative tools like CBCT. The increasing demand for three-dimensional (3D) imaging has propelled CBCT to the forefront of diagnostic techniques in dentistry [28]. This widespread adoption is attributed to several factors, including CBCT's ability to generate high-resolution volumetric images of the jawbone at a cost-effective radiation dose [29]. Additionally, its compact size, affordability, and in-office or close-by availability enhance its clinical utility [30]. Nasal morphology exhibits significant variations across ethnicities [2]. Previous research has explored methods for predicting nasal projection and the position of the pronasale (the most anterior point of the soft tissue nose). These studies have established a link between nasal morphology and the relative positioning of the pronasale, subnasale (the deepest point where the columella meets the upper lip), and nasal alae (outermost points of the nostrils) on the un-

derlying nasal cavity [2, 31].

Limited research has explored the potential of CBCT images to predict nasal position through analysis of the 3D spatial relationships between the underlying skeletal structures and the overlying soft tissues [32]. This study utilized CBCT images to perform measurements of both the soft tissue nose and the corresponding hard tissue nasal aperture. The results demonstrated significant sex-based differences in nasal length, depth, and tip angle. These findings align with previous observations reported by Wang *et al.* [13], Prasad *et al.* [33], and Zamani *et al.* [34].

Previous research conducted on Indonesian and Scottish populations suggests that the nasal index, a ratio of nasal width to height, serves as a reliable anthropometric parameter for geographic origin estimation [29, 35]. Additionally, the nasal index has established utility in forensic science due to its demonstration of sexual dimorphism, aiding in human identification [36]. While our study incorporated nasal index calculations, the results did not reveal statistically significant sex-based differences in this sample. It is important to acknowledge that craniometric parameters, including the nasal index, have been explored in prior research for sex estimation within forensic contexts [15]. However, the validity of these parameters hinges on the understanding that they can vary considerably across populations [37].

Extensive literature highlights significant variations in orbital and nasal morphology influenced by age, sex, and ethnicity [38]. These observations resonate with the findings of Vidya *et al.* [39] who examined South Indian skulls and Nasir *et al.* [40] who investigated nasal indexes across various Indian states. Their studies reported a higher nasal index in males compared to females, which is not necessarily reflected in our specific sample.

The nasolabial angle remains a valuable clinical and cephalometric parameter, playing a crucial role in assessing the soft tissue profile [41]. Research suggests population-based variations in NLA values, with Asian populations generally exhibiting higher values indicative of flatter facial profiles and more obtuse angles compared to Caucasian or African populations [42].

Our study's findings align with this trend, demonstrating significant sex-based differences in the nasolabial angle. These observations are consistent with Kandhasamy *et al.*, who reported higher NLA values in males compared to females [43]. The underlying cause for this sexual dimorphism is likely attributable to variations in the location of the nasolabial angle's vertex point. Kandhasamy *et al.* defined the vertex as the midpoint of tangents drawn along both the columella and the upper lip. This approach positioned their vertex considerably posterior to the posterior columella point, potentially influencing the measured NLA value.

Limitations of the Study

It is important to acknowledge that the current study focused on an Iranian population sample. Nasal morphology exhibits significant variations across ethnicities, limiting the generalizability of these findings to other populations [44].

Despite this limitation, the study offers valuable data on the relationships between the skull and facial soft tissues in the context of nasal reconstruction, along with providing prediction guidelines for this process. These findings hold relevance for applications in forensic science and anthropology. Looking forward, future research should expand its scope

to investigate nasal morphology in other major ethnicities, potentially including populations within Iran. Additionally, employing larger sample sizes would be beneficial in establishing standardized norms for nasal characteristics. These efforts can ultimately contribute to improved accuracy in facial reconstruction techniques and enhanced success rates in forensic identification.

Conclusion

This study revealed significant sex-based variations in nasal parameters. Males exhibited greater nasal length and depth compared to females. Additionally, a notable difference was observed in the nasal tip angle between the sexes, with males having a more projected nasal tip angle. The clinical significance of this study is that hard and soft tissue relation data from CBCT can be useful for predicting the position of the nose. The values of the nasal form are useful for facial reconstruction and rhinoplasty.

Conflict of Interest

The authors have no competing interests to declare that are relevant to the content of this article.

References

1. Xi J, Si XA, Malvè M. Nasal anatomy and sniffing in respiration and olfaction of wild and domestic animals. *Frontiers in Veterinary Science*. 2023 Jul 14;10:1172140.
2. Jankowska A, Janiszewska-Olszowska J, Grocholewicz K. Nasal Morphology and Its Correlation to Craniofacial Morphology in Lateral Cephalometric Analysis. *Int J Environ Res Public Health*. 2021 Mar 16;18(6):3064.
3. Perry JL, Kollara L, Kuehn DP, Sutton BP, Fang X. Examining age, sex, and race characteristics of velopharyngeal structures in 4-to 9-year-old children using magnetic resonance imaging. *The Cleft Palate-Craniofacial Journal*. 2018 Jan;55(1):21-34.
4. Buzek A, Serwańska-Leja K, Zaworska-Zakrzewska A, Kasprzowicz-Potocka M. The shape of the nasal cavity and adaptations to sniffing in the dog (*Canis familiaris*) compared to other domesticated mammals: A review article. *Animals*. 2022 Feb 19;12(4):517.
5. Stenner M, Rudack C. Diseases of the nose and paranasal sinuses in child. *GMS Curr Top Otorhinolaryngol Head Neck Surg*. 2014 Dec 1;13:Doc10.
6. Losco L, Bolletta A, Pierazzi DM, Spadoni D, Cuomo R, Marcasciano M, et al. Reconstruction of the Nose: Management of Nasal Cutaneous Defects According to Aesthetic Subunit and Defect Size A Review. *Medicina (Kaunas)*. 2020 Nov 25;56(12):639.
7. Gupta S, Gupta V, Vij H, Vij R, Tyagi N. Forensic Facial Reconstruction: The

- Final Frontier. *J Clin Diagn Res*. 2015 Sep;9(9):ZE26-8.
8. Damas S, Córdón O, Ibáñez O, Damas S, Córdón O, Ibáñez O. Relationships between the skull and the face for forensic Craniofacial Superimposition. *Handbook on Craniofacial Superimposition: The MEPROCS Project*. 2020:11-50.
9. Li J, Wu S, Mei L, Wen J, Marra J, Lei L, Li H. Facial asymmetry of the hard and soft tissues in skeletal Class I, II, and III patients. *Scientific Reports*. 2024 Feb 29;14(1):4966.
10. Supmaneenukul Y, Khemla C, Parakonthun K. Soft tissue compensation evaluation in patients with facial asymmetry using cone-beam computed tomography combined with 3D facial photographs. *Heliyon*. 2024 Mar 30;10(6):e27720.
11. Kaliappan A, Motwani R, Gupta T, Chandrupatla M. Innovative Cadaver Preservation Techniques: a Systematic Review. *Maedica (Bucur)*. 2023 Mar;18(1):127-135.
12. Thakur S, Sehrawat JS. Age and sex dependent differences in midline facial soft tissue thicknesses measured on MRI scans of Northwest Indian subjects: a forensic anthropological study. *Egyptian Journal of Forensic Sciences*. 2023 Aug 18;13(1):38.
13. Wang J, Wusiman P, Mi C. Cone-beam computed tomography analysis of the nasal morphology among Uyghur nationality adults in Xinjiang for forensic reconstruction. *Translational Research in Anatomy*. 2021 Nov 1;25:100139.
14. Utsuno H, Kageyama T, Uchida K, Kibayashi K, Sakurada K, Uemura K. Pilot study to establish a nasal tip prediction method from unknown human skeletal remains for facial reconstruction and skull photo superimposition as applied to a Japanese male populations. *Journal of Forensic and Legal Medicine*. 2016 Feb 1;38:75-80.
15. Rynn C, Wilkinson CM, Peters HL. Prediction of nasal morphology from the skull. *Forensic Sci Med Pathol*. 2010 Mar;6(1):20-34.
16. Bulut O, Liu CY, Gurcan S, Hekimoglu B. Prediction of nasal morphology in facial reconstruction: Validation and recalibration of the Rynn method. *Legal Medicine*. 2019 Sep 1;40:26-31.
17. Aljabaa AH. Lateral cephalometric analysis of the nasal morphology among Saudi adults. *Clinical, Cosmetic and Investigational Dentistry*. 2019 Jan 15:9-17.
18. Sforza C, Rosati R, De Menezes M, Dolci C, Ferrario VF. Three-dimensional computerized anthropometry of the nose. In *Handbook of Anthropometry: Physical Measures of Human Form in Health and Disease* 2012 Jan 12 (pp. 927-942). New York, NY: Springer New York.
19. Starck WJ, Epker BN. Cephalometric analysis of profile nasal esthetics Part I Method and normative data. *The International Journal of Adult Orthodontics and Orthognathic Surgery*. 1996 Jan 1;11(2):91-103.
20. Jahanbin A, Poosti M, Rashed R, Sharifi V, Bozorgnia Y. Evaluation of nasomaxillary growth of adolescent boys in northeastern Iran. *Acta Medica Iranica*. 2012:684-8.
21. Gruszka K, Aksoy S, Różyło-Kalinowska I, Gülbeş MM, Kalinowski P, Orhan K. A comparative study of paranasal sinus and nasal cavity anatomic variations between the Polish and Turkish Cypriot Population with CBCT. *Head Face Med*. 2022;18(1):1-10.
22. Lee WJ, Wilkinson CM, Hwang HS. An accuracy assessment of forensic computerized facial reconstruction employing cone-beam computed tomography from live subjects. *J Forensic Sci*. 2012;57(2):318-27.
23. Fernandes CMS, de Sena Pereira FDA, da Silva JVL, da Costa Serra M. Is characterizing the digital forensic facial reconstruction with hair necessary A familiar assessors' analysis. *Forensic Sci Int*. 2013;229(1-3):164-73.
24. Wilkinson C. Facial reconstruction— anatomical art or artistic anatomy? *J Anat*. 2010;216(2):235-50.
25. Mala PZ. Pronasale position: an appraisal of two recently proposed methods for predicting nasal projection in facial reconstruction. *Journal of Forensic Sciences*. 2013 Jul;58(4):957-63.
26. Maltais Lapointe G, Lynnerup N, Hoppa RD. Validation of the new interpretation of Gerasimov's nasal projection method for forensic facial approximation using CT data. *Journal of forensic sciences*. 2016 Jan;61:S193-200.
27. Chen F, Chen Y, Yu Y, Qiang Y, Liu M, Fulton D, Chen T. Age and sex related measurement of craniofacial soft tissue thickness and nasal profile in the Chinese population. *Forensic science international*. 2011 Oct 10;212(1-3):272-e1.
28. Bromberg N, Brizuela M. Dental Cone Beam

- Computed Tomography. 2023 Apr 19. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan–.
29. Suomalainen A, Pakbaznejad Esmaceli E, Robinson S. Dentomaxillofacial imaging with panoramic views and cone beam CT. *Insights into imaging*. 2015 Feb;6:1-6.
 30. Sarilita E, Rynn C, Mossey PA, Black S, Oscandar F. Nose profile morphology and accuracy study of nose profile estimation method in Scottish subadult and Indonesian adult populations. *International Journal of Legal Medicine*. 2018 May;132:923-31.
 31. Ryu JY, Park KS, Kim MJ, Yun JS, Lee UY, Lee SS, Roh BY, Seo JU, Choi CU, Lee WJ. Craniofacial anthropometric investigation of relationships between the nose and nasal aperture using 3D computed tomography of Korean subjects. *Sci Rep*. 2020 Sep 30;10(1):16077.
 32. Lee KM, Lee WJ, Cho JH, Hwang HS. Three-dimensional prediction of the nose for facial reconstruction using cone-beam computed tomography. *Forensic science international*. 2014 Mar 1;236:194-e1.
 33. Prasad M, Chaitanya N, Reddy KPK, Talapaneni AK, Myla VB, Shetty SK. Evaluation of nasal morphology in predicting vertical and sagittal maxillary skeletal discrepancies'. *Eur J Dent*. 2014;8(02):197-204.
 34. Zamani Naser A, Panahi Boroujeni M. CBCT evaluation of bony nasal pyramid dimensions in Iranian population: a comparative study with ethnic groups. *Int Sch Res*. 2014;2014(12):25-35.
 35. Sharma SK, Jehan M, Sharma RL, Saxena S, Trivedi A, Bhadkaria V. Anthropometric Comparison of nasal parameters between male and female of Gwalior region. *Journal of Dental and Medical Sciences*. 2014 May;1(5):57-62.
 36. Marini MI, Angrosidy H, Kurniawan A, Margaretha MS. The anthropological analysis of the nasal morphology of Dayak Kenyah population in Indonesia as a basic data for forensic identification. *Translational Research in Anatomy*. 2020 Jun 1;19:100064.
 37. Scendoni R, Kelmendi J, Arrais Ribeiro IL, Cingolani M, De Micco F, Cameriere R. Anthropometric analysis of orbital and nasal parameters for sexual dimorphism: New anatomical evidences in the field of personal identification through a retrospective observational study. *PLoS One*. 2023 May 3;18(5):e0284219.
 38. Khani H, Fazelinejad Z, Hanafi MG, Mahdianrad A, Moghadam AR. Morphometric and volumetric evaluations of orbit using three-dimensional computed tomography in southwestern Iranian population. *Translational Research in Anatomy*. 2023 Mar 1;30:100233.
 39. Vidya CS, Shamasundar NM, Manjunatha B, Raichurkar K. Evaluation of size and volume of maxillary sinus to determine sex by 3D computerized tomography scan method using dry skulls of South Indian origin. *International Journal of Current Research and Review*. 2013; 5: 97.
 40. Nasir N, Asad MR, Muzammil K, Hassan A, Alshalwi T, Khan MR. Anthropometric study of nasal indices in four Indian Stat. *Clin Pract*. 2021; 18: 1620–1625.
 41. Quinzi V, Paskay LC, D'Andrea N, Albani A, Monaco A, Saccomanno S. Evaluation of the nasolabial angle in orthodontic diagnosis: a systematic review. *Applied Sciences*. 2021 Mar 12;11(6):2531.
 42. Rho NK, Park JY, Youn CS, Lee SK, Kim HS. Early changes in facial profile following structured filler rhinoplasty: an anthropometric analysis using a 3-dimensional imaging system. *Dermatologic Surgery*. 2017 Feb 1;43(2):255-63.
 43. Kandhasamy K, Prabu NM, Sivanmalai S, Prabu PS, Philip A, Chiramel JC. Evaluation of the nasolabial angle of the Komarapalayam population. *J Pharm Bioallied Sci*. 2012 Aug;4(Suppl 2):S313-5.
 44. Shah R, Frank-Ito DO. The role of normal nasal morphological variations from race and gender differences on respiratory physiology. *Respir Physiol Neurobiol*. 2022 Mar;297:103823.