

ORIGINAL
ARTICLE

Received 2024-08-28

Revised 2024-09-24

Accepted 2024-11-03

In Vitro Synergistic Antibacterial Effects of Extract and Honey Derived from *Nigella Sativa* on *Streptococcus Mutans*

Seyed-Ali Kamali ¹, Mohammad-Bagher Rezvani ¹, Maryam Pourhajibagher ², Fatemeh Farzaneh ¹,
Hanie-Sadat Emami-Razavi ¹ ✉

¹ Department of Operative Dentistry, Faculty of Dentistry, Shahed University, Tehran, Iran

² Dental Research Center, Dentistry Research Institute, Tehran University of Medical Sciences, Tehran, Iran

Abstract

Background: This study aimed to assess the synergistic antibacterial effects of *Nigella sativa* extract and honey on *Streptococcus mutans*. **Materials and Methods:** In this in vitro study, the minimum inhibitory concentrations (MICs) and minimum bactericidal concentrations (MBCs) of extract and honey derived from *N. sativa* were determined. The antibacterial activity of 1%, 2%, 3%, 4% and 5% extract and honey derived from *N. sativa* was then evaluated against *S. mutans* by the agar disc diffusion method. Data were analyzed by ANOVA followed by pairwise comparisons by the Tukey's test and paired sample t-test. **Results:** *N. sativa* extract showed inhibitory effects on *S. mutans*. The MIC was 250 mg/mL for 1% and 2% concentrations and 125 mg/mL for 3%, 4%, and 5% concentrations of the mixture against *S. mutans*. In addition, the MBC was determined to be 500 mg/mL for the 1% and 2% concentrations, while it was 250 mg/mL for the 3%, 4%, and 5% concentrations. However, *N. sativa* honey did not have any antibacterial activity against *S. mutans*. Combination of *N. sativa* extract and honey had a significant inhibitory effect on *S. mutans*. Amount of 1000 mg/mL concentration of 5% honey and *N. sativa* caused the largest growth inhibition zone (13 mm) with no significant difference with 0.2% chlorhexidine as control (12 mm) ($P=0.74$). **Conclusion:** *N. sativa* honey alone did not demonstrate any significant effect on inhibiting the growth of *S. mutans*. The combination of *N. sativa* and honey had an inhibitory effect on the growth of *S. mutans*. Increased concentration of *N. sativa* result in a greater inhibition of *S. mutans* growth.

[GMJ.2024;13:e3567] DOI: [10.31661/gmj.v13iSP1.3567](https://doi.org/10.31661/gmj.v13iSP1.3567)

Keywords: *Nigella Sativa*; *Streptococcus Mutans*; Honey; Herbal Medicine

Introduction

Dental caries is the most common non-communicable disease, affecting 80% to 90% of the world's population. According to the World Health Organization, approximately half of the world's population (approximately 3.6 billion people) had dental

caries in 2016 [1, 2]. Dental caries is a costly disease, accounting for 5% to 10% of the healthcare budget of industrial countries [3]. Acidogenic and aciduric bacteria such as *Streptococcus mutans* (*S. mutans*), and *Lactobacillus* are responsible for caries development. *S. mutans* has the most prominent role in this regard and it can be abundantly isolated

GMJ

Copyright© 2024, Galen Medical Journal.
This is an open-access article distributed
under the terms of the Creative Commons
Attribution 4.0 International License
(<http://creativecommons.org/licenses/by/4.0/>)
Email: gmj@salviapub.com



✉ Correspondence to:

Hanie-Sadat Emami-Razavi, Department of Operative
Dentistry, Faculty of Dentistry, Shahed University, Teh-
ran, Iran.
Telephone Number: 09124392993
Email Address: Hanie_emami@yahoo.com

ed from carious lesions of humans and animals consuming high-carbohydrate foods [3]. Foods rich in carbohydrates and acidogenic strains of *S. mutans* were long regarded as the main etiology of dental caries. However, recent investigations on *S. mutans* DNA and RNA revealed that it is capable of creating a biofilm and resist the effects of antibiotics [2, 4].

Since *S. mutans* is the main cause of dental caries, the majority of caries control strategies have focused on this microorganism [5, 6].

Some other bacteria may also play a role in development of caries such as *Leptotrichia*, *Rothia*, and *Veillonella* that are involved in the occurrence of enamel caries, and *Atopobium*, *Pseudoramibacter*, *Schlegelella*, *Lactobacillus* and *Streptococcus sanguinis* that are involved in development of dentin caries [7]. Inhibition of *S. mutans* can inhibit caries progression [8, 9].

Nigella sativa (*N. sativa*) from the family of Ranunculaceae is a miracle herb with a long history in traditional medicine. It is native to Southern Europe, North Africa, and southwest Asia, and is cultivated in many Mediterranean and Middle Eastern countries [10]. It has a wide range of therapeutic effects with diuretic, anti-hypertension, anti-diabetic, anti-cancer, immunomodulatory, analgesic, antimicrobial, anti-parasite, anti-inflammation, anti-spasmodic, bronchodilator, and antioxidant properties [11, 12].

Evidence shows that *N. sativa* has inhibitory effects on *S. mutans* and can prevent caries as such [13, 14].

Commercial synthetic mouthwashes such as chlorhexidine and Listerine are commonly used to decrease the oral microbial load [15, 16].

However, due to fewer side effects and higher cost-effectiveness of herbal extracts, researchers have focused on their possible incorporation in the formulation of mouthwashes, given that their safety and optimal antibacterial activity against cariogenic microorganisms are confirmed. *N. sativa* extract has four main alkaloid constituents. It contains thymoquinone, thymol, thymohydroquinone, and di-thymoquinone as effective substances of its aqueous extract [17-19]. Also, *p*-cymene is a major constituent of its extract, which accounts for

higher weight percentage of its volatile oil as well [19]. *N. sativa* contains lipids, vitamins, minerals, proteins (8 essential amino-acids) and carbohydrates [20, 21].

It is also a rich source of essential and non-essential fatty acids. Its main unsaturated fatty acids include linoleic and oleic acids. It also contains phospholipids, carotene, calcium, iron, and potassium [22, 23]. It has antioxidant, anti-inflammatory, anti-histaminic, and immunosupportive effects as well. It can lower the blood glucose and lipid levels, blood pressure, and uric acid [24].

The antibiotics that are effective against *S. mutans* such as ampicillin, cephazolin, methicillin, and clindamycin have side effects such as nausea and vomiting, metallic taste, joint pain, deglutition pain, and stomach burning sensation [25, 26]. Thus, alcoholic extract of *N. sativa* has been considered as an alternative to antibiotics due to its confirmed antibacterial effects [27, 28].

Honey, despite its sweetness, has an inhibitory effect on a number of Gram-positive and Gram-negative bacteria and particularly *S. mutans* [15, 16]. However, its antibacterial activity varies depending on its location of harvesting, type of bees, and honey composition, which may vary from one geographical region to another [18].

Although numerous studies have evaluated the antibacterial activity of *N. sativa* and honey on *S. mutans*, their synergistic antibacterial effect has not been previously investigated. Thus, this study aimed to assess the synergistic antibacterial effects of *N. sativa* extract and *N. sativa* honey on *S. mutans*.

Materials and Methods

This in vitro, experimental study was approved by the ethics committee of the School of Dentistry of Shahed University (IR.SHAHED.REC.1400.192)

Obtaining the Extract and Honey

N. sativa seeds were obtained from the Traditional Medicine Department of Tehran University of Medical Sciences. Its ethanolic extract was subsequently obtained in 70% concentration (200 g of *N. sativa* in 1000 mL of ethanol). [13] After 72 hours, the solution was

filtered through a #1 Whatman filter paper with 150 µm pores, and its ethanol was evaporated by using a water bath (Gesellschaft für Labortechnik GmbH, Burgwedel, Germany). The extract was then stored at 4°C. Since the extract was not water soluble, dimethyl sulfoxide was added to it. The *N. sativa* honey was also collected for this study from the bees fed from *N. sativa* flower of Iran's central area (Isfahan). To mix the *N. sativa* extract and honey, 5 different weight percentages of the *N. sativa* extract (1% to 5% w/w), and 95-99% w/w raw honey were used (for instance, 3% *N. sativa* plus 97% honey). To prepare different concentrations of the mixture, it was serially diluted (from ½ to 1/128 v/v), then 8 different concentrations for each mixture was provided.

S. Mutans Culture

Standard-strain *S. mutans* (ATCC35668) was purchased from the Iranian Biological Resource Center. Then were cultured in 5 ml brain heart infusion (BHI) broth (Merck, Germany) at 37°C in the presence of 5% CO₂ [29]. After that a suspension with a concentration of 1.5×10^8 CFU/mL (equal to 0.5 McFarland) was prepared from it.

Determination of Minimum Inhibitory Concentration (MIC)

The MIC values were determined according to the Clinical and Laboratory Standards Institute (CLSI) standard [30]. In brief, 100 µL of BHI broth was added to the wells of a 96-well microplate. Next, 100 µL of each mixture was added to the first well in final concentration of 1 g/mL. Subsequently, 100 µL of the contents of the first well was transferred to the second well. This process was continued until the 10th well, and finally, 100 µL of the contents of the last well was discarded. Next, 100 µL of bacterial suspension containing 1.5×10^6 CFUs/mL was added to each well, and the microplate was incubated at 37°C and 5% CO₂ for 24 hours.

Wells containing 200 µL of BHI broth served as the negative control while wells containing BHI broth and bacteria served as the positive control. The concentration of mixture in the first well that inhibited visible bacterial growth (turbidity) was recorded as the MIC.

Determination of Minimum Bactericidal Concentration (MBC)

According to the CLSI guideline [31] MBC is identified as the lowest concentration of an antimicrobial agent that can decrease the bacterial population by 99.9% after 24 hours. To determine the MBC, after identifying the MIC, the microplate wells containing the MIC and one higher concentration were inoculated onto BHI agar medium (Merck, Germany). The plates were then incubated at 37°C in an environment with 5% CO₂ for 24 hours. The concentration of mixture at which no bacterial growth was observed was designated as the MBC.

Assessment of Antibacterial Activity by the Agar Disc Diffusion Technique

Paper discs with 6 mm diameter were dipped in 20 µL of the *N. sativa* and honey mixture in different concentrations (similar to those used for MIC determination). *S. mutans* suspension with 0.5 McFarland standard concentration was prepared in BHI broth and swab-cultured on BHI agar (Merck, Germany). The discs were placed on the plate surface with 2 cm distance from each other. The plates were then incubated at 37°C and 5% CO₂ for 24 hours. The diameter of the growth inhibition zones was finally measured by a ruler and reported in millimeters. Chlorhexidine at a concentration of 0.2% was used as the positive control group.

Statistical Analysis

All tests were conducted three times. Normal distribution of data was ensured by the Kolmogorov-Smirnov test. Thus, data were analyzed by ANOVA followed by pairwise comparisons by the Tukey's test. Differences in growth inhibition zone diameters were analyzed by paired sample t-test. $P < 0.05$ was considered statistically significant.

Results

MIC and MBC

Table-1 shows the MIC and MBC of the mixture of *N. sativa* and honey in different concentrations against *S. mutans*. As shown, by an increase in concentration, the antimicrobial activity of the mixture increased. As shown,

Table 1. MIC of the Mixture of N. sativa Extract and Honey in Different Concentrations Against S. mutans

Concentration	MIC (mg/mL)	MBC (mg/mL)
1%	250	500
2%	250	500
3%	125	250
4%	125	250
5%	125	250

the MIC was 125 mg/mL for 3%, 4%, and 5% concentrations of the mixture. While it was 250 mg/mL for 1% and 2% concentrations. Additionally, the MBC was found to be 500 mg/mL for the 1% and 2% concentrations, whereas the MBC for the 3%, 4%, and 5% concentrations was 250 mg/mL.

Agar Disc Diffusion Results

No growth inhibition zone was seen around the discs containing different concentrations of 1% honey and N. sativa. However, 1000 mg/mL concentration of 2% and 3% honey and N. sativa created a *S. mutans* growth inhibition zone by 9 mm. Also, 500 and 1000 mg/mL concentrations of 4% honey and N. sativa caused growth inhibition zones with 10 and 8 mm diameter, respectively. Moreover, 500 mg/mL concentration of 5% honey and N. sativa created a 10 mm growth inhibition zone (Table-2); 1000 mg/mL concentration of 5% honey and N. sativa caused the largest growth inhibition zone (13 mm); however, it had no significant difference with 0.2% chlorhexidine as control (12 mm) ($P=0.74$).

Discussion

This study assessed the synergistic antibacterial effects of N. sativa extract and honey on *S. mutans*. Since thymoquinone, which is the effective substance of N. sativa, has toxic effects in high concentrations [31], low percentages of N. sativa were used in the present study. Also, honey was added to it to improve its taste. The results showed that N. sativa extract had inhibitory effects on *S. mutans*; however, N. sativa honey did not show any antibacterial activity. Combination of N. sativa extract and honey had inhibitory effects on *S. mutans*.

Chaieb *et al.* [32] evaluated the antibacterial activity of thymoquinone, and reported that it had greater effects on Gram-positive bacteria. It also impaired bacterial adhesion. Gawron *et al.* [33] assessed the effect of N. sativa extract on methicillin-resistant *Staphylococcus aureus*, and showed that its antimicrobial activity depended on the concentration of thymoquinone, and could have significant bacteriostatic and bactericidal effects. Mouwakeh *et al.* [34] evaluated the effect of N. sativa extract and bioactive materials on *Staphylococcus aureus* and confirmed its optimal antibacterial activity. The above-mentioned two studies evaluated *Staphylococcus aureus*, which is a Gram-positive facultative anaerobe similar to *S. mutans*. Consistent with their results, the present study showed concentration-dependent antibacterial activity of N. sativa extract against *S. mutans*.

The present results revealed the inhibitory effect of the mixture of N. sativa extract and honey on growth of *S. mutans* in agar disc diffusion technique. Similarly, previous studies showed significant antibacterial activity of N. sativa extract against antibiotic-resistant bacterial species [33, 34]. Dalli *et al.* [35] evaluated the molecular composition and antimicrobial activity of N. sativa on drug-resistant bacteria. They evaluated N. sativa from different geographical sources and showed their variable antimicrobial activity. For instance, N. sativa collected from India had the highest inhibitory effect on *Escherichia coli* followed by N. sativa collected from Saudi Arabia, Morocco and Syria. However, determination of their MIC revealed that all N. sativa types had an inhibitory effect on *Escherichia coli* in concentrations equal or higher than 20 mL. They also showed the optimal antibacterial ef-

Table 2. Diameter of *S. Mutans* Growth Inhibition Zones Caused by Different Concentrations of Honey and *N. Sativa* Extract Mixture

Concentration of mixture	Growth inhibition zone diameter (mm)
1%	-
2% of first concentration	9
3% of first concentration	9
4% of first concentration	10
4% of second concentration	8
5% of first concentration	13
5% of second concentration	10
Negative control (honey)	-
Positive control (chlorhexidine 0.2%)	12

fect of *N. sativa* on *Pseudomonas aeruginosa* and *Acinetobacterbaumannii*[35]. Rostinawat *et al.* [13] showed the anti-*S. mutans* activity of the ethanolic extract of *N. sativa* in 400 to 1200 mg/mL concentrations. The diameter of the growth inhibition zone increased by an increase in concentration of *N. sativa*. Assessment of MIC revealed no bacterial growth in presence of over 38 mg/mL concentrations. Mohammed [36] evaluated the effect of ethanolic and ether extracts of *N. sativa* on *S. mutans* and *Streptococcus mitis*, and reported superior results for the ethanolic extract with a larger growth inhibition zone against *S. mutans*. However, it had lower effects on *Streptococcus mitis*; nonetheless, its ethanolic extract was more effective than its ether extract [36]. Dhahir Mansour Al Sultani *et al.* [37] evaluated the MIC of the lipid and aqueous extracts of *N. sativa* on Gram-positive and Gram-negative bacteria and found their comparable antibacterial activity with no significant difference. Gram-negative bacteria were affected by double the concentration of extract compared with Gram-positive bacteria, indicating higher sensitivity of Gram-positive bacteria to the extract. In the present study, promising results were obtained regarding the antibacterial activity of the extract such that its MIC was 250 mg/mL for 1% and 2% concentrations and 125 mg/mL for 3%, 4%, and

5% concentrations. Also, its 5% concentration showed superior results to chlorhexidine control group in agar disc diffusion test. These results supported the findings of previous studies regarding optimal antibacterial activity of *N. sativa* [35-37]. Azimi Laysar *et al.* [38] showed optimal antibacterial effects of different concentrations of *N. sativa* extract and nano-silver on *S. mutans* and *Streptococcus sanguinis*. The growth inhibition zone caused by nano-silver was larger than that caused by *N. sativa*; nonetheless, the growth inhibition zone caused by *N. sativa* was still larger than that caused by amoxicillin.

In the present study, *N. sativa* honey had no significant inhibitory effect on *S. mutans*, and did not cause its proliferation (due to its sugar content) either. However, this result was in contrast to the findings of previous studies. Rezvani *et al.* [16] confirmed the synergistic effect of honey and cinnamon against *S. mutans*, and reported that this effect was not concentration-dependent. These inconsistent results can be caused by the difference in the composition of cinnamon with *N. sativa* or the difference in the synergistic effect of honey with each of the plant extract. The Nassar *et al.* [39] evaluated the effect of honey on *S. mutans* and biofilm formation. They evaluated both natural and synthetic honey, and showed that bacterial biofilm was easily detached from the wells that contained natural honey while the biofilm firmly adhered to the walls in wells containing synthetic honey. Ahmadi-Motamayelet *et al.* [40] assessed the in vitro effect of honey in different concentrations on *S. mutans* and showed that honey in concentrations >20% had an inhibitory effect on *S. mutans*. It also inhibited *Lactobacillus* in 100% concentration. Deglovic *et al.* [41] demonstrated that honey prevented bacterial adhesion to tooth surface and could be even more effective than CHX. The antibacterial agents in the composition of honey include hydrogen peroxide, which is effective on both Gram-positive and Gram-negative bacteria, gluconic acid, which is effective against Gram-negative bacteria, and defensin-1 which is effective against both Gram-positive and Gram-negative bacteria [41].

Patelet *et al.* [28] demonstrated that honey in concentrations <60% could not inhibit bacte-

rial growth but had optimal antibacterial efficacy in higher concentrations. The growth inhibition zone diameter increased by an increase in its concentration. Roselena *et al.* [42] indicated an increase in antibacterial activity of honey against *S. mutans* by an increase in its concentration. Libonatti *et al.* [43] discussed that honeys obtained from different geographical areas had variable effects on the bacteria. For example, Iranian honey was effective on *Staphylococcus aureus*, and *Pseudomonas aeruginosa* while Malaysian honey was effective on *Salmonella typhi* and *Shigellasonnei*. Voidarou *et al.* [44] evaluated the effects of honeys obtained from different herbs on different bacteria, and showed that thyme honey was effective on *Staphylococcus aureus* with a growth inhibition zone of 4 mm and on *Escherichiacoli* with a growth inhibition zone of 6 mm; whereas, multi-herb honey had no significant effect on *Staphylococcus aureus*, and caused a 3 mm growth inhibition zone in *Escherichia coli* culture. Variations in the results of studies can be due to different origins of honeys.

This study had an in vitro design. Thus, the results can not be directly generalized to the clinical setting. Future studies are required to assess the effects of higher concentrations of *N. sativa* extract and find its safest and most effective concentration for possible incorporation in oral hygiene products. Also, the effects of *N. sativa* extract and honey on expression of colonization and biofilm formation genes of *S. mutans* should be investigated.

Conclusion

N. sativa honey alone did not demonstrate any significant effect on inhibiting the growth of *S. mutans*. The combination of *N. sativa* and honey had an inhibitory effect on the growth of *S. mutans*. Increased concentration of *N. sativa* result in a greater inhibition of *S. mutans* growth.

Conflict of Interest

The authors declare that they have no conflicts of interest related to this article.

References

1. Collaborators GB, To G. Population and fertility by age and sex for 195 countries and territories, 1950–2017. a systematic analysis for the Global Burden of Disease Study. 2017;392(10159):1995-2051.
2. Simón-Soro A, Mira A. Solving the etiology of dental caries. Trends in microbiology. 2015 Feb;23(2):76-82.
3. Moynihan P. Sugars and Dental Caries: Evidence for Setting a Recommended Threshold for Intake. Advances in nutrition (Bethesda, Md.). 2016 Jan 15;7(1):149-56.
4. Petersen PE. Challenges to improvement of oral health in the 21st century--the approach of the WHO Global Oral Health Programme. International dental journal. 2004 Dec;54(6 Suppl 1):329-43.
5. Kt S, Kmk M, Balachander N, Jimson S, Sarangarajan R. Dental caries vaccine-a possible option?. Journal of clinical and diagnostic research: JCDR. 2013 Jun;7(6):1250-3.
6. Mantzourani M, Fenlon M, Beighton D. Association between Bifidobacteriaceae and the clinical severity of root caries lesions. Oral microbiology and immunology. 2009 Feb;24(1):32-7.
7. Lim SM, Lee TK, Kim EJ, Park JH, Lee Y, Bae KS, Kum KY. Microbial profile of asymptomatic and symptomatic teeth with primary endodontic infections by pyrosequencing. Journal of Korean Academy of Conservative Dentistry. 2011 Nov 1;36(6):498-505.
8. Simon L. The Role of Streptococcus mutans And Oral Ecology in The Formation of Dental Caries. Lethbridge Undergraduate Research Journal. 2007;2(2):1718-8482.
9. Acevedo AM, Ray MV, Socorro M, Rojas-Sánchez F. Frequency and distribution of Mutans Streptococci in dental plaque from caries-free and caries-affected Venezuelan children. Acta odontologica latinoamericana: AOL. 2009;22(1):15-20.
10. Khare CP. Encyclopedia of indian medicinal plants: rational western therapy, ayurvedic and other traditional usage, botany. Springer. 2004; : 330-331.
11. Abel-Salam BK. Immunomodulatory effects of black seeds and garlic on alloxan-induced Diabetes in albino rat. Allergologia et immunopathologia. 2012;40(6):336-40.

12. Jahromy MH, Jalili M, Mohajer AJ, Poor FK, Dara SM. Effects of Nigella sativa Seed Extract on Perphenazine-Induced Muscle Rigidity in Male Mice. *World. 2014;4(4):313-8.*
13. Rostinawati T, Karipaya S, Iskandar Y. Antibacterial Activity of Ethanol Extract of Nigella Sativa L Seed Against Streptococcus mutans In IOP Conference Series. *Earth and Environmental Science. 2019;334(1):012050.*
14. Gasong BT, Hartanti AW, Tjandrawinata RR. Antibacterial activity of Nigella sativa L seed oil in water emulsion against dental cariogenic bacteria. *Int J Pharm Sci Res. 2017;8(7):3155-61.*
15. Fernandez MD, Guedes MI, Langa GP, Rösing CK, Cavagni J, Muniz FW. Virucidal efficacy of chlorhexidine: a systematic review. *Odontology. 2021;110:1-7.*
16. Rezvani MB, Niakan M, Kamalinejad M, Ahmadi FS, Hamze F. The synergistic effect of honey and cinnamon against Streptococcus mutans bacteria. *Asian Pacific Journal of Tropical Biomedicine. 2017 Apr 1;7(4):314-20.*
17. Ramadan MF. Nutritional value, functional properties and nutraceutical applications of black cumin (Nigella sativa L.): an overview. *International journal of food science & technology. 2007 Oct;42(10):1208-18.*
18. Sharma NK, Ahirwar D, Jhade D, Gupta S. Medicinal and pharmacological potential of nigella sativa: a review. *Ethnobotanical Leaflets. 2009;13(7):946-55.*
19. Geng D, Zhang S, Lan J. Analysis on chemical components of volatile oil and determination of thymoquinone from seed of Nigella glandulifera. *China journal of Chinese materia medica. 2009 Nov;34(22):2887-90.*
20. Ghosheh OA, Houdi AA, Crooks PA. High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed (Nigella sativa L.). *Journal of pharmaceutical and biomedical analysis. 1999 Apr;19(5):757-62.*
21. Al-Jassir MS. Chemical composition and microflora of black cumin (Nigella sativa L.) seeds growing in Saudi Arabia. *Food Chemistry. 1992 Jan 1;45(4):239-42.*
22. Nickavar B, Mojab F, Javidnia K, Amoli MA. Chemical composition of the fixed and volatile oils of Nigella sativa L from Iran *Zeitschrift für Naturforschung. C, Journal of Biosciences. 2003 Sep 1;58(9-10):629-31.*
23. Ramadan MF, Mörsel JT. Characterization of phospholipid composition of black cumin (Nigella sativa L.) seed oil. *Die Nahrung. 2002 Aug;46(4):240-4.*
24. Ali BH, Blunden G. Pharmacological and toxicological properties of Nigella sativa. *Phytotherapy Research: PTR. 2003 Apr 1;17(4):299-305.*
25. Al-Shami IZ, Al-Hamzi MA, Al-Shamahy HA, Majeed AL. Efficacy of some Antibiotics against Streptococcus Mutans Associated with Tooth decay in Children and their Mothers. *On J Dent and Oral Health. 2019;2(1):1-4.*
26. Little WA, Thomson LA, Bowen WH. Antibiotic susceptibility of Streptococcus mutans: comparison of serotype profiles. *Antimicrobial agents and chemotherapy. 1979 Mar;15(3):440-3.*
27. Jebali N, Rabbani Khorasgani M, Kianfar M, Emami H. Evaluation of the effects of honey, vinegar and rosewater on adhesion and biofilm formation of streptococcus mutans and streptococcus Sobrinus. *Journal of Isfahan Dental School. 2016;12(3):232-47.*
28. Patel HR, Krishnan CA, Thanveer K. Antimicrobial effect of honey on Streptococcus mutans—An in vitro study. *International Journal of Dental Science and Research. 2013;2(1):46-9.*
29. Liu J, Wu C, Huang IH, Merritt J, Qi F. Differential response of Streptococcus mutans towards friend and foe in mixed-species cultures. *Microbiology. 2011 Sep;157(9):2433-44.*
30. Performance standards for antimicrobial susceptibility testing; Twenty-second Informational supplement. In: CLSI document M100-S22. Wayne, PA: Clinical and Laboratory Standards Institute. 2020;32:m100.
31. Abukhader MM. The effect of route of administration in thymoquinone toxicity in male and female rats. *Indian Journal of Pharmaceutical Sciences. 2012 May 1;74(3):195-200.*
32. Chaieb K, Kouidhi B, Jrah H, Mahdouani K, Bakhrouf A. Antibacterial activity of Thymoquinone, an active principle of Nigella sativa and its potency to prevent bacterial biofilm formation. *BMC complementary and alternative medicine. 2011 Apr 13;11:29.*
33. Gawron G, Krzyckiowski W, Lemke K, Ołdak A, Kadziński L, Banecki B. Nigella sativa seed extract applicability in preparations against methicillin-resistant

- Staphylococcus aureus and effects on human dermal fibroblasts viability. Journal of ethnopharmacology. 2019 Nov 15;244:112135.
34. Mouwakeh A, Kincses A, Nové M, Mosolygó T, Mohácsi-Farkas C, Kiskó G, Spengler G. Nigella sativa essential oil and its bioactive compounds as resistance modifiers against Staphylococcus aureus. Phytotherapy research: PTR. 2019 Apr;33(4):1010-8.
35. Dalli M, Azizi SE, Benouda H, Azghar A, Tahri M, Bouammali B, Maleb A, Gseyra N. Molecular Composition and Antibacterial Effect of Five Essential Oils Extracted from Nigella sativa L. Seeds against Multidrug-Resistant Bacteria: A Comparative Study. Evidence-based complementary and alternative medicine: eCAM. 2021 Mar 15;2021:6643765.
36. Mohammed NA. Effect of Nigella Sativa L extracts against Streptococcus mutans and Streptococcus mitis in Vitro. J Bagh College Dentistry. 2012;24(3):154-7.
37. Dhahir Mansour Al Sultani T, AyadFakhri S, Al-Zuhairi WH. Comparison of the Antimicrobial Activity of Nigella sativa Aqueous and Oil Extracts. Iranian Journal of War and Public Health. 2021 Sep 10;13(4):305-11.
38. AzimiLaysar H, Niakan M, Mohammad Taghi G, Jafarian Z, Mostafavizade M, Niakan S. Comparison of the antibacterial activity of various concentrations of Nigella Sativa and Nanosilver on the growth of S sanguis and S mutans. Res Dent Sci. 2013 Jan 10;9(4):179-86.
39. Nassar HM, Li M, Gregory RL. Effect of honey on Streptococcus mutans growth and biofilm formation. Applied and environmental microbiology. 2012 Jan;78(2):536-40.
40. Ahmadi-Motamayel F, Hendi SS, Alikhani MY, Khamverdi Z. Antibacterial activity of honey on cariogenic bacteria. Journal of dentistry (Tehran, Iran). 2013 Jan;10(1):10-5.
41. Deglovic J, Majtanova N, Majtan J. Antibacterial and Antibiofilm Effect of Honey in the Prevention of Dental Caries: A Recent Perspective. Foods (Basel, Switzerland). 2022 Sep 2;11(17):2670.
42. Roselena B, Mayra C, Juana S. Antibacterial effect of honey in different concentrations on Streptococcus mutans Oral Session Cariology. Cariogenic Plaque Inhibitors. 2012 Jun 23; 12: .
43. Libonatti C, Varela S, Basualdo M. Antibacterial activity of honey: A review of honey around the world. Journal of Microbiology and Antimicrobials. 2014 Mar 31;6(3):51-6.
44. Voidarou C, Alexopoulos A, Plessas S, Karapanou A, Mantzourani I, Stavropoulou E, Fotou K, Tzora A, Skoufos I, Bezirtzoglou E. Antibacterial activity of different honeys against pathogenic bacteria. Anaerobe. 2011 Dec;17(6):375-9ZhongguoZhongyaozhizhi=Zhongguozhongyaozhizhi=China journal of Chinese materia medica. 2009 Nov;34(22):2887-90.