

ORIGINAL
ARTICLE

Received 2024-10-29

Revised 2024-12-09

Accepted 2024-12-21

Investigating The Antifungal Activity of Atorvastatin Compared with Nystatin on Oral Candidiasis before and during Head and Neck Radiotherapy

Zahra Golestannejad¹, Faezeh Khozeimeh¹, Sara Taheri², Soha Feizi³, Fatemeh Abbasi⁴, Zahra Saberi¹, Elham Faghiehian¹, Parvin Dehghan⁵, Faezeh Tadayon⁶, Mahnaz Kheirkhah⁷, Zeinab Nourmohammadi Najafabadi⁸, Zahra Ataie⁹✉

¹ Department of Oral and Maxillofacial Medicine, Dental Research Center, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

² School of Dentistry, Dental Research Center, Dental Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

³ Department of Stomatology, Xi'an Jiaotong University, Xi'an, China

⁴ Department of Oral Medicine, Dental Research Center, Dental Research Institute, Faculty of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

⁵ Department of Mycology and Parasitology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

⁶ Department of Dentistry, Dental Students Research Committee, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

⁷ Department of Mycology and Parasitology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

⁸ Dentistry Department, Isfahan University of Medical Science, Isfahan, Iran

⁹ Dental Research Center, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: Radiotherapy plays an imperative role in the control of head and neck malignancies; however, it can damage oral tissues and salivary glands. These damages can alter oral *Candida* species and lead to the expansion of oral candidiasis resistant to common antifungal mediators, including nystatin. Atorvastatin, a serum cholesterol-lowering drug, has potential antifungal activities by inhibiting the synthesis of ergosterol in the fungal wall and disrupting mitochondrial function. This study aimed to determine and equivalence the antifungal activities of nystatin and atorvastatin on *Candida* species isolated from the mouths of patients undergoing head and neck radiotherapy before and during radiotherapy. **Materials and Methods:** This was an in vitro laboratory research conducted on samples isolated from patients experiencing head and neck radiotherapy, before and during radiotherapy. After determining the *Candida* species using the PCR-RFLP method, the antifungal activity of both nystatin and atorvastatin was evaluated by microdilution method according to CLSI standards, and the minimum inhibitory concentration (MIC) and minimum lethal concentration (MFC) of each drug were measured. **Results:** According to our findings, atorvastatin had less activity in inhibiting and killing different *Candida* species compared to nystatin before and during radiotherapy. Before radiotherapy, the MIC and MFC indices for nystatin against *Candida albicans* ($P < 0.001$), tropical ($P < 0.001$) and glabrata ($P < 0.001$) were significantly lower than these three indices ($P\text{-value} < 0.001$) for atorvastatin. The results of these indices in the second week of radiotherapy were similar to the results before radiotherapy. The MIC and MFC results for *Candida albicans* and *tropicalis* were obtained with a $P\text{-value} < 0.001$ and for glabrata with a $P\text{-value} = 0.002$. **Conclusion:** The discoveries of this study indicate that atorvastatin exhibits lower antifungal activity compared to nystatin in treating oral candidiasis among patients receiving head and neck radiotherapy, both before and during the treatment. [GMJ.2024;13:e3717] DOI: [10.31661/gmj.v13iSP1.3717](https://doi.org/10.31661/gmj.v13iSP1.3717)

Keywords: Oral Candidiasis; Antifungal Activity; Nystatin; Atorvastatin; Radiotherapy

GMJ

Copyright© 2024, Galen Medical Journal.
This is an open-access article distributed
under the terms of the Creative Commons
Attribution 4.0 International License
(<http://creativecommons.org/licenses/by/4.0/>)
Email: gmj@salviapub.com



✉ Correspondence to:

Zahra Ataie, Dental Research Center, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran.

Telephone Number: 009831-37925502

Email Address: zataei@gmail.com

Introduction

One of the effective modalities in controlling head and neck malignancies is radiotherapy. Most of those suffering from these types of malignancies require radiotherapy as primary treatment, after surgery, in combination with chemotherapy, or as palliative treatment. Although radiotherapy has known antitumor activity, ionizing radiation can also damage healthy tissues in the treated area [1-2]. Head and neck radiotherapy often causes damage to oral and facial structures, including the major salivary glands [3] and can lead to complications such as inflammation of the oral mucosa (mucositis), qualitative and quantitative changes in saliva, and dry mouth, about one to two weeks after the initiation of the treatment, which in turn predisposes patient to oral fungal proliferation, colonization, and infections [4-7].

Dry mouth is one of the symptoms that indicate a decrease in salivary secretion due to radiotherapy of the salivary glands. Atrophy of salivary gland tissue can reduce the quality and especially the quantity of saliva, causing an acidification of salivary pH, a decrease in antimicrobial capacity and a decrease in the cleansing effect of the mouth. All these salivary changes affect oral normal microbial flora, especially *Candida* species; facilitate the presence of pathogenic agents and increase the risk of oral candidiasis infection [5, 8].

Head and neck radiotherapy not only increases the number of fungal species, but also changes their strains [2, 9, 10]. Candidiasis is the furthest common clinical infection of the oropharynx in patients undergoing radiotherapy [11]. Studies since 1990 have shown that in addition to *Candida albicans*, non-*albicans* species also play a role in the colonization and infection of patients undergoing radiotherapy [6, 7, 9]. *Candida albicans* remains the predominant species in oral infections; however, non-*albicans* species are also emerging as important pathogens, and infections caused by the simultaneous presence of these species are more severe and more resistant to treatment [5, 6, 12].

Oral candidiasis infections can cause unpleasant complications such as burning sensation in the mouth, pain and dysphagia, which may

lead to temporary or complete interruption of radiotherapy, and reduces the antitumor effects of treatment [5]. Moreover, during severe immunosuppression, oral candidiasis can spread to deeper organs and be fatal. Therefore, rapid diagnosis and proper management and treatment of this infection are essential [8]. Among the routine antifungal drugs currently used to control oral candidiasis, topically and systemically; nystatin is a polyene that directly binds to ergosterol in the cell membrane of fungi, leading to the formation of pores in the membrane, disruption of ion balance and cell death. However, the changing epidemiology of fungal infections and the emergence of strains resistant to conventional treatments, including nystatin, have created new challenges in the treatment of these infections [13, 14].

Statins, which have been used as cholesterol-lowering agents in recent decades, have multiple effects, including antifungal activity [15]. By inhibiting the HMGCR enzyme, which is involved in the production of ergosterol in fungi, as in humans; these drugs can alter the fluidity and lipid composition of fungal membranes and disrupt cellular metabolism [15-18].

Considering the inhibitory activity of atorvastatin on fungal growth and the necessity of new drugs in the face of treatment-resistant strains, present experimental survey objectives to determine the frequency of different oral *Candida* species in patients with head and neck malignancies before and during radiotherapy and to investigate the sensitivity of these species to nystatin and atorvastatin.

Materials and Methods

Study Samples

This investigation was conducted as an experimental study (in vitro). The *Candida* species examined were previously collected and preserved in the Department of Parasitology and Mycology at Isfahan University of Medical Sciences from 33 patients diagnosed with head and neck malignancies at Seyed al-Shohaday Hospital in Isfahan, before and during their radiotherapy treatment. Ethical approval for the research was granted by Isfahan University of Medical Sciences, and informed

consent was obtained from all participants before the study commenced. Initially, the cohort comprised 18 women and 15 men, aged between 38 and 74 years, with 21 individuals exhibiting *Candida* infections prior to the initiation of radiotherapy. During the second week of treatment, 6 patients (four men and two women) succumbed, and 19 out of the remaining 27 patients developed *Candida* infections. Additionally, some patients were found to be infected with multiple strains of *Candida* concurrently, such as *Candida albicans* and *Candida tropicalis*. The isolated strains were identified and characterized using the restriction fragment length polymorphism PCR (PCR-RFLP) technique. The *Candida* strains were initially cultured on sub-dextrose agar (SDA) medium and incubated at 35°C for 24 hours to prepare a suspension. Laboratory procedure

To prepare the suspension of the candida, a portion of the *Candida* species was added to 1 ml of distilled water and the optical absorption of the resulting suspension was measured at a wavelength of 530 nm using a WPA Bio-wave II spectrophotometer (Biochrom UK). The addition of the species to the suspension continued until its optical absorption reached the range of 0.13-0.08. At this stage, the cell concentration of the resulting suspension was equivalent to 0.5 McFarland. In order to prepare the initial stock of the two antifungal drugs under study, 12.5 mg of nystatin powder (Sigma-Aldrich, Germany) and 12.5 mg of atorvastatin powder (Merck, Germany) were dissolved in 1 ml of methanol and 12.5 mg of atorvastatin powder (Merck, Germany) were dissolved in 1 ml of dimethyl sulfoxide or DMSO (Merck-Germany), respectively; Affording to the Clinical and Laboratory Standards Institute (CLSI) standard; and reserved at laboratory temperature for 30 minutes until the resulting stock was homogenized.

Drug susceptibility testing

To assess the antifungal efficacy of nystatin and atorvastatin, separate evaluations of the minimum inhibitory concentration (MIC) and minimum lethal concentration (MFC) were performed at 24 and 48 hours for various *Candida* species, including *Candida albicans*, *Candida glabrata*, *Candida tropicalis*, *Candida*

parapsilosis, and *Candida krusei*. The MIC for both agents was established through a serial dilution technique utilizing 96-well ELISA microplates. For nystatin, ten wells for concentrations of 0.5-128 µg/ml and for atorvastatin, ten wells for concentrations of 8-1024 µg/ml, as well as two positive and negative control wells were prepared. Subsequently, 100 µl of fungal suspension from each species was introduced into each of the ten wells. In the remaining two wells, 100 µl of organism suspension at a density of 1×10^3 cells/ml was combined with 100 µl of Roswell Park Memorial Institute (RPMI) medium in the positive control well, while 100 µl of RPMI containing the drug was added to the negative control well, in accordance with the CLSI-M27 guidelines. Following incubation at 35°C for 24 and 48 hours, the turbidity in the wells was assessed, with the first well exhibiting no turbidity identified as the minimum inhibitory concentration (MIC). This process allowed for the determination of MIC₂₄ and MIC₄₈.

To ascertain the minimum lethal concentration (MFC), 100 µl of the suspension from the MIC well and the subsequent wells were transferred to Sabouraud Dextrose Agar (SDA) medium. After performing a sweep culture and incubating for 48 hours at 35°C, the plate that displayed five or fewer colonies of the target species was designated as the MFC. To determine the break point for nystatin, $2 \mu\text{g/ml} < \text{MIC}_{48}$ was considered as drug resistance and $2 \mu\text{g/ml} > \text{MIC}_{48}$ was considered as drug susceptibility. For atorvastatin, since it is not essentially an antifungal drug, a break point is not identified.

Data Analysis

The data were analyzed using SPSS software package, version 22 (SPSS Inc., Chicago, Ill., USA). To investigate the antifungal activity of both nystatin and atorvastatin, three indices MIC₂₄, MIC₄₈ and MFC of each of them were calculated separately on *Candida albicans*, *Candida glabrata* and *Candida tropicalis* species and their median, range and mode were determined. To compare the antifungal activity of these two drugs, the values of the aforementioned indices before and during radiotherapy intervention were analyzed using Mann-Whitney statistical test.

Results

The *Candida* samples studied from the fungal collection of the Department of Parasitology and Mycology, Isfahan University of Medical Sciences, included 14 *Candida albicans*, 5 *Candida tropicalis*, and 2 *Candida glabrata* samples before the start of radiotherapy, and 12 *Candida albicans*, 4 *Candida tropicalis*, 2 *Candida glabrata*, 1 *Candida parapsilosis*, and 1 *Candida krusei* sample in the second week of radiotherapy.

Tables-1 and 2 revealed the median and range of MIC₂₄, MIC₄₈, and MFC indices of the two drugs, nystatin and atorvastatin, against *Candida albicans*, *glabrata*, and *tropicalis*. Antifungal indices were also determined for *Parapsilosis* and *Krusei* species, but due to

the small number of samples, comparing their frequency distribution was not possible. Also, considering these two species were not isolated from any samples before radiotherapy, comparison of antifungal indices of nystatin with atorvastatin before and during radiotherapy for these two species was not plausible. Based on the findings, MIC₂₄, MIC₄₈, and MFC indices of nystatin against *Candida parapsilosis* were 0.5, 0.5, and 1 µg/ml, respectively. The MIC₂₄, MIC₄₈, and MFC of atorvastatin were 32, 64, and 128 µg/ml against *Candida parapsilosis* and 64, 256, and 256 µg/ml against *Candida krusei*, respectively.

To compare the antifungal activity of nystatin against atorvastatin, the levels of three indices (MIC₂₄, MIC₄₈, and MFC) of these two drugs were separately analyzed by Mann-Whitney

Table 1. Median and range of the MIC₂₄, MIC₄₈, and MFC indices of nystatin on strains isolated from the patients before and during radiotherapy intervention

Indices		<i>Candida albicans</i>		<i>Candida tropicalis</i>		<i>Candida glabrata</i>	
		Before RT	During RT	Before RT	During RT	Before RT	During RT
Antifungal activity							
MIC ₂₄	Median	1	0.75	1	1	1	0.5
	Range	<0.2-5	0.1-5	0.1-5	0.1-5	1-1	0.5-0.5
MIC ₄₈	Median	1	1.5	2	1.5	2	0.5
	Range	0.4-5	0.2-5	0.2-5	2-1	2-1	0.5-0.5
MFC	Median	1	1	1	1	1	0.5
	Range	0.32-5	<0.1-0.5	0.1-5	0.8-5	0.2-5	0.5-0.5

Table 2. Median and range of the MIC₂₄, MIC₄₈ and MFC indices of atorvastatin on strains isolated from patients before and during radiotherapy intervention

Indices		<i>Candida albicans</i>		<i>Candida tropicalis</i>		<i>Candida glabrata</i>	
		Before RT	During RT	Before RT	During RT	Before RT	During RT
Antifungal activity							
MIC ₂₄	Median	128	128	128	128	128	384
	Range	256 - 64	256 - 128	512 - 32	512 - 64	128 - 64	512 - 256
MIC ₄₈	Median	256	256	256	384	256	384
	Range	1024 - <128	1024 - <256	1024 - <64	1024 - <256	1024 - <256	512 - 256
MFC	Median	512	768	1024	1024	1025	768
	Range	1024 - <128	1024 - <128	1024 - <128	1024 - <512	1024 - <1024	1024 - 512

once before treatment and once during radiotherapy treatment on each species, and the results are listed in Tables-3 and 4. Comparison of the antifungal activity of nystatin and atorvastatin showed that the MIC₂₄, MIC₄₈, and MFC indices for nystatin both before and during radiotherapy were significantly lower than atorvastatin ($P < 0.001$). These values were obtained for *Candida albicans* and tropicalis species with $P < 0.001$ and *Candida glabrata* species with $P = 0.002$.

Discussion

The findings of the current investigation indicated that prior to and throughout the course of radiotherapy, nystatin exhibited a superior antifungal efficacy compared to atorvastatin against all *Candida* species obtained from patients receiving head and neck radiotherapy. Specifically, all three antifungal efficacy metrics—MIC₂₄, MIC₄₈, and MFC—were found to be lower for nystatin in comparison to atorvastatin. Atorvastatin's antifungal effects are primarily mediated through the inhibition of HMG-CoA reductase, which reduces ergosterol synthesis—a critical component of fungal cell membranes. Unlike nystatin, which directly binds to ergosterol and disrupts membrane integrity, atorvastatin's effect is in-

direct and less potent at achievable therapeutic concentrations. Recent studies underscore this limitation, noting that while atorvastatin exhibits inhibitory activity against a range of *Candida* species and azole-resistant strains, its efficacy significantly lags behind nystatin due to lower bioavailability and potential resistance mechanisms, such as alternative sterol synthesis pathways. Furthermore, atorvastatin's ability to disrupt fungal mitochondria offers an avenue for future exploration, particularly in combination therapies with azoles like fluconazole, which have shown enhanced efficacy against azole-resistant strains. [19, 20]

Lima [21] and Nyilasi [22] compared the antifungal activity of nystatin against atorvastatin on *Candida* species. In the Lima study, which was conducted on fungal strains from the American Cell Bank cultured in mice, the MIC of atorvastatin was lower for *Candida albicans* and *Candida glabrata* species and higher for nystatin than current study.

In *Candida tropicalis*, unlike the previous two species, the MIC of atorvastatin in the Lima study was higher (256 $\mu\text{g/ml}$) than the present study (128 $\mu\text{g/ml}$); however, in both studies, the MIC of nystatin activity on *Candida tropicalis* was almost equivalent.

The study of Nyilasi [22] showed similar results to the present study regarding the anti-

Table 3. Comparison of antifungal indices (MIC₂₄, MIC₄₈, and MFC) of nystatin against atorvastatin, based on *Candida* species isolated from patients undergoing radiotherapy, before radiotherapy intervention

Indices		<i>Candida albicans</i>		<i>Candida tropicalis</i>		<i>Candida glabrata</i>	
		Nystatin	Atorvastatin	Nystatin	Atorvastatin	Nystatin	Atorvastatin
Antifungal Activity							
MIC ₂₄	Median	1	128	1	128	1	128
	Range	<0.2 - 5	256 - 64	0.15	512 - 32	1 - 1	128 - 64
	P-value	<0.001		<0.001		<0.001	
MIC ₄₈	Median	1	256	2	256	2	256
	Range	0.4 - 5	1024 - 128	0.2 - 5	1024 - <128	2 - 1	11024 - <256
	P-value	<0.001		<0.001		<0.001	
MFC	Median	1	512	1	1024	1	1025
	Range	0.32 - 5	1024 - <128	0.1 - 5	1024 - <128	0.2 - 5	1024 - 1024<
	P-value	<0.001		<0.001		<0.001	

Table 4. Comparison of antifungal indices (MIC24, MIC48, and MFC) of nystatin against atorvastatin, based on candida species isolated from patients undergoing radiotherapy, during radiotherapy intervention

Indices		<i>Candida albicans</i>		<i>Candida tropicalis</i>		<i>Candida glabrata</i>	
		Nystatin	Atorvastatin	Nystatin	Atorvastatin	Nystatin	Atorvastatin
Antifungal Activity	Median	0.75	128	1	128	0.5	384
	Range	0.1 - 5	256 - 128	0.1 - 5	512 - 64	0.5 - 0.5	512 - 256
	P-value	<0.001		<0.001		0.002	
MIC48	Median	1.5	256	1.5	384	0.5	384
	Range	0.2 - 5	1024 - <256	2 - 1	1024 - <256	0.5 - 0.5	512 - 256
	P-value	<0.001		<0.001		0.002	
MFC	Median	1	768	1	1024	0.5	768
	Range	1 - <0.05	1024 - <128	0.8 - 5	1024 - <512	0.5 - 0.5	1024 - 512
	P-value	<0.001		<0.001		0.002	

fungus activity of nystatin and atorvastatin on *Candida albicans* species, so the MIC of atorvastatin and nystatin on *Candida albicans* species in both studies was 128 µg/ml and 1 µg/ml, respectively. However, the results for *Candida glabrata* species were different regarding atorvastatin drug and were 32 µg/ml in Nyilasi study and 128 µg/ml in the present study; moreover, the results, the MIC of atorvastatin for *Candida glabrata* species increased to 328 µg/ml after radiotherapy.

These differences could be due to the difference in the source of the studied species and also the difference in the environment of the isolated *Candida* species. So in the present study, the change in the quantity and quality of saliva after radiotherapy may lead to a change in the pathogenesis and virulence of the species [23].

The Brillhante study [24] showed that atorvastatin has very different efficacy against different *Candida* species, with the mean MIC values of 52.6 µg/ml for *Candida albicans*, 165.34 µg/ml for *Candida tropicalis*, 755.06 µg/ml for *Candida crusei*, and 1491.37 µg/ml for *Candida parapsilosis*.

The results of current study showed that, contrary to the Brillhante's findings, the MIC for the parapsilosis species was lower than the other studied species and much lower than the results obtained in the Brillhante study

(MIC for the parapsilosis species in the present study and in the Brillhante study: 32 µg/ml - 1491.37 µg/ml, respectively). Similar to the parapsilosis species, the *Candida crusei* species also had a lower MIC in the present study compared to the Brillhante study (MIC for the *crusei* species in the present study and the Brillhante study: 256 µg/ml and 755 µg/ml, respectively). However, for the *Candida albicans* and *tropicalis*, the results of the present study were consistent with the results of his study (MIC for *albicans* in the present study and the Brillhante study were 128 µg/ml and 6.52 µg/ml, respectively, and MIC for *tropicalis* in the present study and the Brillhante study were 128 µg/ml and 165 µg/ml, respectively). The difference in the source of the species and the differences in the method used to measure the MIC, including the solvent of atorvastatin in these studies can explain the differences in the results of the current and aforementioned studies. Ting's systematic study showed that the MIC of statins changes under the influence of factors such as the choice of solvent type, statin form used, methodology and culture medium, and exposure time to statin [25]. In the present study, the solvent type was DMSO, the statin form was pure powder (Pure, Fine, Powdered), the microdilution method, SDA culture medium, and the exposure time was 24 and 48 hours. However, in the Brillhante

study, the solvent was sterile distilled water, and the culture medium and statin form were not specified.

Atorvastatin has been widely used as a serum cholesterol-lowering drug since 1990 [26] and has recently been studied as a novel antifungal drug with an inhibitory effect on fungal walls' biosynthesis [27], so that it may be a suitable alternative for the control of oral candidiasis in cases where the potential for drug resistance is likely. Radiotherapy, by changing the oral environmental conditions including xerostomia, changes in saliva quality, changes in oral pH and mucosal damage (mucositis), leads to changes in the physiological and morphological behavior of *Candida* species and ultimately changes in the pathogenesis and virulence of yeast [6, 23, 27, 28].

Karbach et al. (2012) shown that decreased salivation following head and neck radiotherapy was associated with the emergence of treatment-resistant *Candida* strains [29]. Research indicates that alterations in environmental conditions can prompt swift modifications in the gene expression of *Candida albicans*, facilitating its adaptation to these changes [30]. Paula and Zida determined that 3% of HIV-positive individuals without clinical signs of candidiasis and 4.8% of symptomatic hospitalized patients exhibited resistance to nystatin among various *Candida* species [31, 32]. In contrast to this perspective, the findings of the current study revealed that all strains obtained from patients receiving radiotherapy demonstrated sensitivity to nystatin (MIC < 2 µg/ml) both before and following treatment. This observation aligns with the results reported by Bulacio [33] and Kurnatowski [34].

Furthermore, atorvastatin, even at the maximum concentration tested (1024 µg/ml), still allowed for fungal growth in several instances. Another important hypothesis is that, given the lack of gastrointestinal absorption of nystatin and its only local efficacy, cases of oral candidiasis in people undergoing radiotherapy, who may be immunocompromised and develop candidemia due to factors such as underlying malignancy, previous or concurrent chemotherapy with radiotherapy, or mucositis and subsequent malnutrition [4], require antifungal drugs with systemic and not only

local effects. Atorvastatin has gastrointestinal absorption and systemic effects and may be used as a suitable drug in the aforementioned cases where candidemia is likely to occur. The serum level of atorvastatin in Iranians with a daily dose of 40 mg has been reported to be 50 ± 30 ng/ml [35-37]. The results showed that the minimum concentration of atorvastatin that can have an inhibitory effect on *Candida* is 32 µg/ml. Given this concentration is approximately 1000 times the serum level of atorvastatin, this drug cannot be prescribed to control possible candidemia following orofacial candidal infection in individuals undergoing radiotherapy.

Furthermore, the obtained results showed that the antifungal activity of the two drugs (inhibitory effect-killing effect) is not impacted by radiotherapy. While Ramla showed that the production of hydrolytic enzymes including phospholipase and proteinase by the fungus can increase following radiotherapy [6], which itself may change the efficacy of antifungal drugs following radiotherapy. However, since these enzymes are contributed with the fungus ability to colonize at the host tissue and invade the tissue, they do not affect the susceptibility of the species to antifungal drugs in vitro.

While this study indicates that atorvastatin is less effective than nystatin in treating oral candidiasis during radiotherapy, it still holds potential in specific clinical scenarios. For systemic candidiasis, atorvastatin's gastrointestinal absorption and systemic effects may offer advantages, particularly in immunocompromised patients where localized treatments like nystatin are insufficient.

It is recommended that the antifungal activity of these two drugs compared in vivo in the further studies and to investigate the synergistic effect of atorvastatin when added to the nystatin therapy regimen, rather than comparing the clinical effect of atorvastatin against nystatin.

This study has several limitations. The sample size was relatively small, particularly for certain *Candida* species such as *Candida glabrata*, *Candida parapsilosis*, and *Candida krusei*, which may limit how widely the results can be applied, and a larger group of participants would provide more reliable

Conclusion

Overall, the results showed that atorvastatin is not a appropriate alternative to nystatin in patients undergoing radiotherapy.

The antifungal activity of atorvastatin, both before and during radiotherapy, was less than

that of nystatin in the treatment of oral candidiasis in patients experiencing head and neck radiotherapy.

Conflict of Interest

None.

References

- Vissink A, Jansma J, Spijkervet F, Burlage F, Coppes R. Oral sequelae of head and neck radiotherapy. *Crit Rev Oral Biol Med*. 2003;14(3):199-212.
- Golestannejad Z, Khozeimeh F, Dehghan P, Najafizadeh N, Faghihian E, Kheirkhah M, et al. Comparison of the antifungal effect of voriconazole and fluconazole on oral candidiasis before and during radiotherapy. *DRJ*. 2022;19:99.
- Leung WK, Dassanayake RS, Yau JY, Jin LJ, Yam WC, Samaranyake LP. Oral colonization, phenotypic, and genotypic profiles of *Candida* species in irradiated, dentate, xerostomic nasopharyngeal carcinoma survivors. *J Clin Microbiol*. 2000;38(6):2219-26.
- Schelenz S, Abdallah S, Gray G, Stubbings H, Gow I, Baker P, et al. Epidemiology of oral yeast colonization and infection in patients with hematological malignancies, head neck and solid tumors. *J Oral Pathol Med*. 2011;40(1):83-9.
- Mañas A, Cerezo L, de la Torre A, García M, Albuquerque H, Ludeña B, et al. Epidemiology and prevalence of oropharyngeal candidiasis in Spanish patients with head and neck tumors undergoing radiotherapy treatment alone or in combination with chemotherapy. *Clin Transl Oncol*. 2012;14(10):740-6.
- Ramla S, Sharma V, Patel M. Influence of cancer treatment on the *Candida albicans* isolated from the oral cavities of cancer patients. *Supportive care in cancer: Support Care Cancer*. 2016;24(6):2429-36.
- Jham BC, França EC, Oliveira RR, Santos VR, Kowalski LP, da Silva Freire AR. *Candida* oral colonization and infection in Brazilian patients undergoing head and neck radiotherapy: a pilot study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;103(3):355-8.
- Suryawanshi H, Ganvir SM, Hazarey VK, Wanjare VS. Oropharyngeal candidosis relative frequency in radiotherapy patient for head and neck cancer. *JOMFP*. 2012;16(1):31.
- Paula CR, Sampaio MCC, Birman EG, Siqueira AM. Oral yeasts in patients with cancer of the mouth, before and during radiotherapy. *Mycopathologia*. 1990;112(2):119-24.
- Torres SR, Peixoto CB, Caldas DM, Silva EB, Akiti T, Nucci M, et al. Relationship between salivary flow rates and *Candida* counts in subjects with xerostomia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2002;93(2):149-54.
- Epstein JB, Freilich MM, Le ND. Risk factors for oropharyngeal candidiasis in patients who receive radiation therapy for malignant conditions of the head and neck. *Oral Surg Oral Med Oral Pathol*. 1993;76(2):169-74.
- Redding SW. The role of yeasts other than *Candida albicans* in oropharyngeal candidiasis. *Curr Opin Infect Dis*. 2001;14(6):673-7.
- Prasad R, Shah AH, Rawal MK. Antifungals: Mechanism of Action and Drug Resistance. *Adv Exp Med Biol*. 2016;892:327-49.
- Rajni E, Chaudhary P, Garg VK, Sharma R, Malik M. A complete clinico-epidemiological and microbiological profile of candidemia cases in a tertiary-care hospital in Western India. *ASHE*. 2022;2(1):e37.
- Westermeyer C, Macreadie IG. Simvastatin reduces ergosterol levels, inhibits growth and causes loss of mtDNA in *Candida glabrata*. *FEMS Yeast Res*. 2007;7(3):436-41.
- Nyilasi I, Kocsube S, Pesti M, Lukács G, Papp T, Vágvölgyi C. In vitro interactions between primycin and different statins in their effects against some clinically important fungi. *J Med Microbiol*. 2010;59(2):200-5.
- Westermeyer C, Macreadie IG. Simvastatin reduces ergosterol levels, inhibits growth and causes loss of mtDNA in *Candida glabrata*. *FEMS Yeast Res*. 2007;7(3):436-41.
- Katirae F, Teifoori F, Soltani M. Emergence of azole-resistant *Candida* species in AIDS patients with oropharyngeal candidiasis in Iran. *Curr Med Mycol*. 2015;1(3):11-6.
- Esfahani AN, Golestannejad Z, Khozeimeh F, Dehghan P, Maheronnaghsh M, Zarei Z. Antifungal effect of Atorvastatin against *Candida* species in comparison to

- Fluconazole and Nystatin. *Med Pharm Rep.* 2019;92(4):368-73.
20. de Oliveira Neto AS, Souza ILA, Amorim MES, de Freitas Souza T, Rocha VN, do Couto RO, et al. Antifungal efficacy of atorvastatin-containing emulgel in the treatment of oral and vulvovaginal candidiasis. *Med Mycol.* 2021;59(5):476-85.
 21. Lima WG, Alves-Nascimento LA, Andrade JT, Vieira L, de Azambuja Ribeiro RIM, Thomé RG, et al. Are the Statins promising antifungal agents against invasive candidiasis? *BIOMED PHARMACOTHER.* 2019;111:270-81.
 22. Nyilasi I. Effect of different statins on the antifungal activity of polyene antimycotics. *Acta Biol Szeged.* 2010;54(1):33-6.
 23. Davis D. Adaptation to environmental pH in *Candida albicans* and its relation to pathogenesis. *Curr Genet.* 2003;44(1):1-7.
 24. Brilhante RSN, Caetano EP, Oliveira JSd, Castelo-Branco DdSC, Souza ERY, Alencar LPd, et al. Simvastatin inhibits planktonic cells and biofilms of *Candida* and *Cryptococcus* species. *BJID.* 2015;19(5):459-65.
 25. Ting M, Whitaker EJ, Albandar JM. Systematic review of the in vitro effects of statins on oral and perioral microorganisms. *Eur J Oral Sci.* 2016;124(1):4-10.
 26. Kontoyiannis DP. Decrease in the number of reported cases of zygomycosis among patients with diabetes mellitus: a hypothesis. *Clin Infect Dis.* 2007;44(8):1089-90.
 27. Sun Y, Cao C, Jia W, Tao L, Guan G, Huang G. pH regulates white-opaque switching and sexual mating in *Candida albicans*. *Eukaryot Cell.* 2015;14(11):1127-34.
 28. Bensadoun RJ, Patton LL, Lalla RV, Epstein JB. Oropharyngeal candidiasis in head and neck cancer patients treated with radiation: update 2011. *Supportive care in cancer: Support Care Cancer.* 2011;19(6):737-44.
 29. Karbach J, Walter C, Al-Nawas B. Evaluation of saliva flow rates, *Candida* colonization and susceptibility of *Candida* strains after head and neck radiation. *Clin Oral Investig.* 2012;16(4):1305-12.
 30. Wilson D, Thewes S, Zakikhany K, Fradin C, Albrecht A, Almeida R, et al. Identifying infection-associated genes of *Candida albicans* in the postgenomic era. *FEMS Yeast Res.* 2009;9(5):688-700.
 31. Zida A, Yacouba A, Bamba S, Sangare I, Sawadogo M, Guiguemde T, et al. In vitro susceptibility of *Candida albicans* clinical isolates to eight antifungal agents in Ouagadougou (Burkina Faso). *JMM.* 2017;27(4):469-75.
 32. de Paula SB, Morey AT, Santos JP, dos Santos PM, Gameiro DG, Kerbauy G, et al. Oral *Candida* colonization in HIV-infected patients in Londrina-PR, Brazil: antifungal susceptibility and virulence factors. *JIDC.* 2015;9(12):1350-9.
 33. Bulacio L, Paz M, Ramadán S, Ramos L, Pairoba C, Sortino M, et al. Oral infections caused by yeasts in patients with head and neck cancer undergoing radiotherapy, Identification of the yeasts and evaluation of their antifungal susceptibility. *Journal de mycologie medicale.* 2012;22(4):348-53.
 34. Kurnatowski P, Moqbil S, Kaczmarczyk D. Signs, symptoms and the prevalence of fungi detected from the oral cavity and pharynx of radiotherapy subjects with head and neck tumors, and their susceptibility to chemotherapeutics. *Ann Parasitol.* 2014;60(3):207-13.
 35. Burlage FR, Coppes RP, Meertens H, Stokman MA, Vissink A. Parotid and submandibular/sublingual salivary flow during high dose radiotherapy. *Radiotherapy and oncology: Radiother Oncol.* 2001;61(3):271-4.
 36. Dambroso D, Svidzinski Tie, Svidzinski Ae, Dalalio M, Machado DO, Moliterno RA. Radiotherapy effect on frequency of *Candida* spp and on virulence of *C albicans* isolated from the oral cavity of head and neck cancer patients. *Rev Ciênc Farm Básica Apl.* 2009;30(2):153-9.
 37. Chou Y-C, Wang Y-K, Charng M-J, Ueng Y-F. Determination of serum atorvastatin concentrations in lipid-controlling patients with and without myalgia syndrome. *JFDA.* 2013;21(2):147-53.